

## Clonetics™ normal human prostate cell systems

### Instructions for use

---

#### Safety statements

**THESE PRODUCTS ARE FOR RESEARCH USE ONLY.** Not approved for human or veterinary use, for application to humans or animals, or for use in clinical or *in vitro* procedures.

**WARNING: CLONETICS™ AND POIETICS™ PRODUCTS CONTAIN HUMAN SOURCE MATERIAL, TREAT AS POTENTIALLY INFECTIOUS.** Each donor is tested and found non-reactive by an FDA approved method for the presence of HIV-1, hepatitis B virus and hepatitis C virus. Where donor testing is not possible, cell products are tested for the presence of viral nucleic acid from HIV, hepatitis B virus, and hepatitis C virus. Testing can not offer complete assurance that HIV-1, hepatitis B virus, and hepatitis C virus are absent. All human sourced products should be handled at the biological safety level 2 to minimize exposure of potentially infectious products, as recommended in the CDC-NIH manual, Biosafety in Microbiological and Biomedical Laboratories, 5<sup>th</sup> edition. If you require further information, please contact your site safety officer or scientific support.

#### Unpacking and storage instructions

1. Check all containers for leakage or breakage.
2. For cryopreserved cells – remove cryovials from the dry ice packaging and immediately place into liquid nitrogen storage. Alternatively, thaw and use the cells immediately. If no dry ice remains, please contact customer service.
3. For proliferating cells – swab down the flask of proliferating cells with 70% ethanol or isopropanol, then place the flask in 37°C, 5% CO<sub>2</sub>, humidified incubator and allow to equilibrate for 3 to 4 hours. After cells have equilibrated, remove shipping medium from the flask and replace with fresh medium.
4. BulletKit™ instructions: upon arrival, store basal medium at 4-8°C and SingleQuots™ at –20°C in a freezer that is not self-defrosting. If thawed upon arrival, growth factors can be stored at 4°C and added to basal medium within 72 hours of receipt. After SingleQuots™ are added to basal medium, use within 1 month. Do not re-freeze.
5. ReagentPack™ subculture reagents are sterile-filtered and then stored at –20°C until shipment. Subculture reagents may thaw during transport. They may be refrozen once. If you plan to use within 3 days, store at 4°C. Trypsin/EDTA solu-

tion has a limited shelf life or activation at 4°C. If, upon arrival, trypsin/EDTA is thawed, immediately aliquot and refreeze at –20°C. We recommend that the HEPES-BSS and the trypsin neutralizing solution be stored at 4°C for no more than 1 month.

**NOTE:** To keep trypsin/EDTA fresh and active after thawing, you may aliquot it into sterile centrifuge tubes and re-freeze at –20°C.

Using media or reagents other than what's recommended will void the cell warranty. Please contact scientific support if you need help selecting media and/or reagents.

#### Preparation of media

**For a BulletKit™, perform the following steps:**

1. Decontaminate the external surfaces of all supplement vials and the medium bottle with ethanol or isopropanol.
2. Aseptically open each supplement vial and add the entire amount to the basal medium with a pipette.
3. Rinse each cryovial with the medium. It may not be possible to recover the entire volume listed for each cryovial. Small losses, even up to 10%, should not affect the cell growth characteristics of the supplemented medium.
4. Transfer the label provided with each kit to the basal medium bottle being supplemented. Use it to record the date and amount of each supplement added. We recommend that you place the completed label over the basal medium label (avoid covering the basal medium lot # and expiration date) to avoid confusion or possible double supplementation.
5. Record the new expiration date on the label based on the shelf life.

**Note:** If there is concern that sterility was compromised during the supplementation process, the entire newly prepared growth medium may be refiltered with a 0.2µm filter to assure sterility. Routine refiltration is not recommended.

#### Thawing of cells / initiation of culture process

1. The recommended seeding density for PrEC is 2,500 cells/cm<sup>2</sup> and the recommended seeding density for PrSC is 3,500 cells/cm<sup>2</sup>.
2. To set up cultures calculate the number of vessels needed based on the recommended seeding density and the surface area of the vessels being used. Do not seed cells into a well plate directly out of cryopreservation. Add the appropriate amount of medium to the vessels (1 ml/5 cm<sup>2</sup>) and allow the vessels to equilibrate in a 37°C, 5% CO<sub>2</sub>, humidified incubator for at least 30 minutes.
3. Wipe cryovial with ethanol or isopropanol before opening. In a sterile field, briefly twist the cap a quarter turn to relieve pressure, then retighten. Quickly thaw the cryovial in a 37°C water bath being careful not to submerge the entire vial. Watch your cryovial closely; when the last sliver of ice melts remove it. Do not submerge it completely. Thawing the cells for longer than 2 minutes results in less than optimal results.
4. Resuspend the cells in the cryovial and using a micropipette, dispense cells into the culture vessels set up earlier. Gently rock the culture vessel to evenly distribute the cells and return to the incubator.
5. Centrifugation should not be performed to remove cells from cryoprotectant cocktail. This action is more damaging than the effects of DMSO residue in the culture.
3. Prepare new culture vessels.
4. Subculture 1 flask at a time. All flasks following the first flask will be subcultured following an optimization of this protocol based on calculated cell count, cell viability, and seeding density.

#### In a sterile field:

1. Aspirate the medium from 1 culture vessel.
2. Rinse the cells with 5 ml of room temperature HEPES-BSS. DO NOT forget this step. The medium contains complex proteins and calcium that neutralize the trypsin.
3. Aspirate the HEPES-BSS from the flask.
4. Cover the cells with 2 ml of trypsin/EDTA solution.
5. Examine the cell layer microscopically.
6. Allow the trypsinization to continue until approximately 90% of the cells are rounded up. This entire process takes about 2-6 minutes, depending on cell type.
7. At this point, rap the flask against the palm of your hand to release the majority of cells from the culture surface. If only a few cells detach, you may not have let them trypsinize long enough. Wait 30 seconds and rap again. If cells still do not detach, wait and rap every 30 seconds thereafter.
8. After cells are released, neutralize the trypsin in the flask with 4 ml of room temperature TNS. If the majority of cells do not detach within 7 minutes, the trypsin is either not warm enough or not active enough to release the cells. Harvest the culture vessel as described above, and either re-trypsinize with fresh, warm trypsin/EDTA solution or rinse with TNS and then add fresh, warm medium to the culture vessel and return to an incubator until fresh trypsinization reagents are available.
9. Quickly transfer the detached cells to a sterile 15 ml centrifuge tube.
10. Rinse the flask with a final 2 ml of HEPES-BSS to collect residual cells, and add this rinse to the centrifuge tube.
11. Examine the harvested flask under the microscope to make sure the harvest was successful by looking at the number of cells left behind. This should be less than 5%.
12. Centrifuge the harvested cells at 220 x g for 5 minutes to pellet the cells.
  - a. Aspirate most of the supernatant, except for 100-200 µl.
  - b. Flick the cryovial with your finger to loosen the pellet.

#### Subculturing

The following instructions are for a 25 cm<sup>2</sup> flask. Adjust all volumes accordingly for other size flasks.

Note: Lonza warrants its Clonetics™ cells only if Lonza subculturing reagents are used. The recommended subculturing reagents for these cells are trypsin/EDTA (CC-5012), trypsin neutralizing solution (CC-5002), and HEPES buffered saline solution (CC-5022). These reagents can be purchased individually or together as part of the ReagentPack™ subculture reagents (CC-5034).

#### Preparation for subculturing the first flask:

1. Subculture the cells when they are 60%-80% confluent and contain many mitotic figures throughout the flask.
2. For each 25 cm<sup>2</sup> of cells to be subcultured:
  - a. Thaw 2 ml of trypsin/EDTA and allow to come to room temperature.
  - b. Allow 7-10 ml of HEPES buffered saline solution (HEPES-BSS) to come to room temperature.
  - c. Allow 4 ml of trypsin neutralizing solution (TNS) to come to room temperature.
  - d. Remove growth medium from 4°C storage and allow to start warming to room temperature.

for a single procedure, transfer and warm only the required volume to a sterile secondary container.

13. Dilute the cells in 2-3 ml of growth medium and note the total volume of the diluted cell suspension.
14. Determine cell count and viability using a hemacytometer and trypan blue. Make a note of your cell yield for later use.
15. If necessary, dilute the suspension with the HEPES-BSS to achieve the desired "cells/ml" and re-count the cells.
16. Use the following equation to determine the total number of viable cells.

$$\text{Total \# of Viable Cells} = \frac{\text{Total cell count} \times \text{percent viability}}{100}$$

17. Determine the total number of flasks to inoculate by using the following equation. The number of flasks needed depends upon cell yield and seeding density. If seeding into well plates at this time, the recommended density is 10,000 cells/cm<sup>2</sup>.

$$\text{Total \# of Flasks to inoculate} = \frac{\text{Total \# of viable cells}}{\text{Growth area} \times \text{Rec. Seeding Density}}$$

18. Use the following equation to calculate the volume of cell suspension to seed into your flasks.

$$\text{Seeding Volume} = \frac{\text{Total volume of diluted cell suspension}}{\text{\# of flasks as determined in step 18}}$$

19. Prepare flasks by labeling each flask with the passage number, strain number, cell type and date.
20. Carefully transfer growth medium to new culture vessels by adding 1 ml growth medium for every 5 cm<sup>2</sup> surface area of the flask (1 ml/5 cm<sup>2</sup>).
21. After mixing the diluted cells with a 5 ml pipet to ensure a uniform suspension, dispense the calculated volume into the prepared subculture flasks.
22. If not using vented caps, loosen caps of flasks. Place the new culture vessels into a 37°C humidified incubator with 5% CO<sub>2</sub>.

## Maintenance

1. Change the growth medium the day after seeding and every other day thereafter. As the cells become more confluent, increase the volume of media as follows: under 25% confluence then feed cells 1 ml per 5 cm<sup>2</sup>, 25-45% confluence then feed cells 1.5 ml per 5 cm<sup>2</sup>, over 45% confluence then feed cells 2 ml per 5 cm<sup>2</sup>.
2. Warm an appropriate amount of medium to 37°C in a sterile container. Remove the medium and replace it with the warmed, fresh medium and return the flask to the incubator.
3. Avoid repeated warming and cooling of the medium. If the entire contents are not needed

All trademarks herein are marks of Lonza Group or its subsidiaries.

## Ordering information

Cryopreserved cells (single donor)		
CC-2555	PrEC	≥500,000 cells
CC-2508	PrSC	
Proliferating cells in flasks		
CC-2655	PrEC	T-25 flask
CC-0310	PrEC	T-75 flask
CC-2608	PrSC	T-25 flask
CC-2508T75	PrSC	T-75 flask
Proliferating cells in plates		
CC-2555W6	PrEC	6-well plate
CC-2555W48	PrEC	48-well plate
CC-0088	PrEC	96-well plate
CC-2508W6	PrSC	6-well plate
CC-2508W48	PrSC	48-well plate
CC-2508W96	PrSC	96-well plate

For additional proliferating formats, please go to [www.lonza.com](http://www.lonza.com).

## Related products

### Prostate epithelial growth media (must be purchased separately):

CC-3166	PrEGM™ BulletKit™	Kit which contains a 500 ml bottle of PrEBM™, (CC-3165) and PrEGM™ SingleQuots™ (CC-4177).
CC-3165	PrEBM™	Prostate epithelial basal medium (no growth factors) (500 ml)
CC-4177	PrEGM™ SingleQuots™	Supplements and growth factors (BPE, hydrocortisone, hEGF, epinephrine, insulin, triiodothyronine, transferrin, gentamicin/amphotericin-B and retinoic acid)

**Stromal cell growth media (must be purchased separately):**

CC- 3205	SCGM™ BulletKit™	Kit which contains a 500 ml bottle of SCBM™, (CC-3204) and SCGM™ SingleQuotes™ (CC-4181).
CC-3204	SCBM™	Stromal cell basal medium (no growth factors) (500 ml)
CC-4181	SCGM™ SingleQuotes™	Supplements and growth factors (hFGF-B, insulin, FBS and gentamicin/ amphotericin-B)

**Product warranty**

CULTURES HAVE A FINITE LIFESPAN *IN VITRO*. Lonza warrants its cells in the following manner only if Clonetics™ media and reagents are used.

1. Clonetics™ PrEC and PrSC cryopreserved cultures are assured for experimental use for 15 population doublings.
2. Clonetics™ PrEC and PrSC proliferating cultures are assured for experimental use for 10 population doublings.
3. Additional population doublings and subcultures are possible, but growth rate, biological responsiveness and function deteriorate with subsequent passage.
4. PrEC and PrSC can become irreversibly contact-inhibited if allowed to reach confluence. To avoid the loss of your cells and forfeiture of your warranty, subculture cells before they reach 80% confluence.

**Quality control**

HIV-1, hepatitis B and hepatitis C are not detected for all donors and/or cell lots. Routine characterization of PrEC includes positive immunofluorescent staining for cytokeratin (clone 8.13). PrSC test positive for vimentin and negative for pan cytokeratin. For detailed information concerning QC testing, please refer to the certificate of analysis.