

## Product References

### Poietics® CD14+ Monocyte Cells

---

1. \*Fujiyama S, Amano K, Uehira K, Yoshida M, Nishiwaki Y, Nozawa Y, Jin D, Takai S, Miyazaki M, Egashira K, Imada T, Iwasake T, Matsubara H. Bone marrow monocyte lineage cells adhere on injured endothelium in a monocytes chemoattractant protein-1-dependent manner and accelerate reendothelialization as endothelial progenitor cells. *Circ Res.* 2003 Nov; 93(10): 980-9.
2. \*Hori S, Heike Y, Takei M, Maruyama M, Inoue Y, Lee JJ, Kim HJ, Harada Y, Kawai H, Shimosaka A, Kami M, Tanosaki MD, Wakasugi H, Saito S, Takaue T, Kakizoe T. Freeze-thawing procedures have no influence on the phenotypic and functional development of dendritic cells generated from peripheral blood CD14+ monocytes. *J Immunother.* 2004; 27(1): 27-35.
3. Kuwana M, Okazaki Y, Kodama H, Izumi K, Yasuoka H, Ogawa Y, kawakami T, Okeda Y. Human circulating CD14+ monocytes as a source of progenitors that exhibit mesenchymal cell differentiation. *J Leukoc Biol.* 2003 Nov; 75(5): 833-45.
4. \*Merck E, Gaillard C, Gorman DM, Montero-Julian F, Durand I, Zurawski SM, Menetrier-Caux C, Carra G, Lebecque S, Trinchieri G, Bates EE. OSCAR is an FcRγ-associated receptor expressed by myeloid cells, involved in antigen presentation and activation of human dendritic cells. *Blood.* 2004; 104(5): 1386-95.
5. \*Nissinen R, Leirisalo-Repo M, Peltomaa Rm Palosuo T, Vaarala O. Cytokine and chemokine receptor profile of peripheral blood mononuclear cells during treatment with infliximab in patients with active rheumatoid arthritis. *Ann Rheum Dis.* 2004 Jun; 63(6): 681-7.
6. Wang C, Jiao C, Hanlon HD, Zheng W, Tomanek RJ, Schatteman GC. Mechanical, cellular, and molecular factors interact to modulate circulating endothelial cell progenitors. *Am J Physiol Heart Circ Physiol.* 2004; 286: H1985-93.

### Macrophage Differentiation

1. \*Chomarat P, Banchereau J, Davoust J, Palucka AK. IL-6 switches the differentiation of monocytes from dendritic cells to macrophages. *Nat Immunol.* 2000 Dec; 1(6): 510-4.
2. \*Kramer JL, Baltathakis I, Alcantara OS, Boldt DH. Differentiation of functional dendritic cells and macrophages from human peripheral blood monocyte precursors is dependent on expression of p21 (WAF1/CIP1) and requires iron. *Br J Haematol.* 2002 Jun; 117(3): 727-34.
3. \*Lee CK, Kim JK, Kim Y, Lee MK, Kim K, Kang JK, Hofmeister R, Durum SK, Han SS. Generation of macrophages from early T progenitors in vitro. *J Immunol.* 2001 May 15; 166(10): 5964-9.
4. \*Lu R, Pitha PM. Monocyte differentiation to macrophage requires interferon regulatory factor 7. *J Biol Chem.* 2001 Nov 30; 276(48): 45491-6.
5. \*Mochida-Nishimura K, Akagawa KS, Rich EA. Interleukin-10 contributes development of macrophage suppressor activities by macrophage colony-stimulating factor, but not by granulocyte-macrophage colony-stimulating factor. *Cell Immunol.* 2001 Nov 25; 214(1): 81-8.
6. \*Olikowsky T, Wang ZQ, Dudhane A, Horowitz H, Conti B, Hoffmann MK. Two distinct pathways of human macrophage differentiation are mediated by interferon-gamma and interleukin-10. *Immunology.* 1997 May; 91(1): 104-8.

7. \*Plesner A, Greenbaum CJ, Lernmark A.  
Low serum conditions for in vitro generation  
of human macrophages with macrophage  
colony stimulating factor. J Immunol  
Methods. 2001 Mar 1; 249(1-2): 53-61.
8. \*Winston BW, Krein PM, Mowat C, Huang  
Y. Cytokine-induced macrophage  
differentiation: a tale of 2 genes. Clin Invest  
Med. 1999 Dec; 22(6): 236-55.

\* References not specifically citing the use of Lonza cells, media,  
or reagents in their research.

+ Denotes sections containing only the articles published within  
the last ten years.