

Product References

Clonetics® Keratinocyte Cells

Cells*

1. Aizu T, Tamai K, Nakano H, Rokunohe D, Toyomaki Y, Uitto J, Sawamura D. Calcineurin/NFAT-dependent regulation of 230-kDa bullous pemphigoid antigen (BPAG1) gene expression in normal human epidermal keratinocytes. *J Dermatol Sci*. 2008 Jul;51(1):45-51.
2. Arlian LG, Morgan MS, Peterson KT. House Dust and Storage Mite Extracts Influence Skin Keratinocyte and Fibroblast Function. *Int Arch Allergy Immunol*. 2008;145(1):33-42.]
3. Bhat KR, Benton BJ, Rosenthal DS, Smulson ME, Ray R. Role of poly(ADP-ribose) polymerase (PARP) in DNA repair in sulfur mustard-exposed normal human epidermal Keratinocytes (NHEK). *J Applied Toxicol*. 2000; 20(S1): S13-7.
4. Bogiatzi SI, Fernandez I, Bichet JC, Marloie-Provost MA, Volpe E, Sastre X, Soumelis V. Cutting Edge: Proinflammatory and Th2 cytokines synergize to induce thymic stromal lymphopoietin production by human skin keratinocytes. *J Immunol*. 2007 Mar 15;178(6):3373-7.
5. Canning MT, Brown DA, Yarosh DB. A bicyclic monoterpene diol and UVB stimulate BRCA1 phosphorylation in human keratinocytes. *Photochem Photobiol*. 2003 Jan; 77(1): 46-51.
6. Cerimele F, Currell F, Ely S, Friedman-Kien AE, Cesarman E, Flore O. Kaposi's sarcoma-associated herpesvirus can productively infect primary human Keratinocytes and alter their growth properties. *J Virol*. 2001 Mar; 75(5): 2435-43.
7. Chan E-C, Chang A, Wu T-Z, Lin Y-L. Effects of recombinant lysostaphin on cytotoxicity and interleukin-8 level in normal human epidermal keratinocytes cell line. *Biotechnol Lett*. 1998 Nov; 20(11): 1053-6.
8. Cheong HI, Johnson J, Cormier M, Hosseini K. In vitro cytotoxicity of eight beta-blockers in human corneal epithelial and retinal pigment epithelial cell lines: Comparison with epidermal keratinocytes and dermal fibroblasts. *Toxicol In Vitro*. 2008 Jun;22(4):1070-6.
9. Ching T-T, Maunakea A, Jun P, Hong C, Zardo G, Pinkel D, Albertson DG, Fridlyang J, Mao J-H, Shchors K, Weiss WA, Costello JF. Epigenome analyses using BAS micorarrays identify evolutionary conservation of tissue-specific methylation of SHANK3. *Nat Genet*. 2005 Jun; 37(6): 645-51.
10. Chou CC, Yang JH, Chen SD, Monteiro-Riviere NA, Li HN, Chen JJ. Expression profiling of human epidermal keratinocyte response following 1-minute JP-8 exposure. *Cutan Ocul Toxicol*. 2006; 25(2):141-53.
11. Cohen EE, Lingen MW, Zhu B, Zhu H, Straza MW, Pierce C, Martin LE, Rosner MR. Protein kinase C zeta mediates epidermal growth factor-induced growth of head and neck tumor cells by regulating mitogen-activated protein kinase. *Cancer Res*. 2006; 66(12):6296-303.
12. Dasgupta S, Tripathi PK, Qin H, Bhattacharya-Chatterjee M, Valentino J, Chatterjee SK. Identification of molecular targets for immunotherapy of patients with head and neck squamous cell carcinoma. *Oral Oncol*. 2006; 42(3): 306-16.
13. Dong S, Kanno T, Yamaki A, Kojima T, Shiraiwa M, Kawada A, Méchin MC, Chavanas S, Serre G, Simon M, Takahara H. NF-Y and Sp1/Sp3 are involved in the transcriptional regulation of the peptidylarginine deiminase type III gene (PADI3) in human keratinocytes. *Biochem J*. 2006; 397(3):449-59.
14. Eckhart L, Schmidt M, Mildner M, Mlitz V, Abtin A, Ballaun C, Fischer H, Mrass P, Tschachler E. Histidase expression in human epidermal keratinocytes: regulation by differentiation status and all-trans retinoic

- acid. *J Dermatol Sci*. 2008 Jun;50(3):209-15.
15. Edlundh-Rose E, Kupershmidt I, Gustafsson AC, Parasassi T, Serafino A, Bracci-Laudiero L, Greco G, Krasnowska EK, Romano M-C, Lundeberg T, Nilsson P, Lundeberg J. Gene expression analysis of human epidermal keratinocytes after N-acetyl L-cysteine treatment demonstrates cell cycle arrest and increased differentiation. *Pathobiol*. 2005; 72(4): 203-12.
 16. Fischer H, Eckhart L, Mildner M, Jaeger K, Buchberger M, Ghannadan M, Tschachler E. DNase1L2 degrades nuclear DNA during corneocyte formation. *J Invest Dermatol*. 2007 Jan;127(1):24-30.
 17. Gabet AS, Accardi R, Bellopede A, Popp S, Boukamp P, Sylla BS, Londoño-Vallejo JA, Tommasino M. Impairment of the telomere/telomerase system and genomic instability are associated with keratinocyte immortalization induced by the skin human papillomavirus type 38. *FASEB J*. 2008 Feb;22(2):622-32.
 18. Gebbia JA, Coleman JL, Benach JL. *Borrelia spirochetes* upregulate release and activation of matrix metalloproteinase gelatinase B (MMP-9) and collagenase 1 (MMP-1) in human cells. *Infect Immun*. 2001 Jan; 69(1): 456-62.
 19. Gibson S, Shillitoe EJ. Analysis of apoptosis-associated genes and pathways in oral cancer cells. *J Oral Pathol Med*. 2006; 35: 146-54.
 20. Gray LJ, Bjelogric P, Appleyard VC, Thompson AM, Jolly CE, Lain S, Herrington CS. Selective induction of apoptosis by leptomycin B in keratinocytes expressing HPV oncogenes. *Int J Cancer*. 2007 Jun 1;120(11):2317-24.
 21. Gross CL, Nealley EW, Nipwoda MT, Smith WJ. Pretreatment of human epidermal keratinocytes with D,L-sulforaphane protects against sulfur mustard cytotoxicity. *Cutan Ocul Toxicol*. 2006;25(3):155-63.
 22. Guo X, Ruiz A, Rando RR, Bok D, Gudas LJ. Esterification of all-trans-retinol in normal human epithelial cell strains and carcinoma lines from oral cavity, skin and breast: reduced expression of lecithin:retinol acyltransferase in carcinoma lines. *Carcinogen*. 2000; 21(11): 1925-33.
 23. Gustafsson AC, Kupershmidt I, Edlundh-Rose E, Greco G, Serafino A, Krasnowska EK, Lundeberg T, Bracci-Laudiero L, Romano M-C, Parasassi T, Lundeberg J. Global gene expression analysis in time series following N-acetyl L-cysteine induced epithelial differentiation of human normal and cancer cells in vitro. *BMC Canc*. 2005 Jul 07; 5: 75-94.
 24. Halder B, Bhattacharya U, Mukhopadhyay S, Giri AK. Molecular mechanism of black tea polyphenols induced apoptosis in human skin cancer cells: involvement of Bax translocation and mitochondria mediated death cascade. *Carcinogenesis*. 2008 Jan;29(1):129-38.
 25. Hasina R, Whipple ME, Martin LE, Kuo WP, Ohno-Machado L, Lingen MW. Angiogenic heterogeneity in head and neck squamous cell carcinoma: biological and therapeutic implications. *Lab Invest*. 2008 Apr;88(4):342-53.
 26. Ho M, Yang E, Marcuk G, Deng D, Sampas N, Tsalenko A, Tabibiazar R, Zhang Y, Chen M, Talbi S, Ho YD, Wang J, Tsao PS, Bendor A, Yakhini Z, Bruhn L, Quertermous T. Identification of endothelial cell genes by combined database mining and microarray analysis. *Physiol Genom*. 2003; 13: 249-62.
 27. Hsu S. A mechanism-based in vitro anticancer drug screening approach for phenolic phytochemicals. 2003 Oct; 1(5): 611-8.
 28. Janssen FP, Bouten CV, van Leeuwen GM, van Steenhoven AA. Effects of temperature and doxorubicin exposure on keratinocyte damage in vitro. *In Vitro Cell Dev Biol Anim*. 2008 Mar-Apr;44(3-4):81-6.
 29. Jin X, Ray R, Leng Y, Ray P. Molecular determination of laminin-5 degradation: a biomarker for mustard gas exposure diagnosis and its mechanism of action. *Exp Dermatol*. 2008 Jan;17(1):49-56.
 30. Jin X, Ray R, Ray P. An immunochromatographic assay to detect reduced level of laminin-5 gamma2 in sulfur mustard-exposed normal human epidermal keratinocytes. *J Appl Toxicol*. 2008 Aug;28(6):759-64.
 31. Kee S-H, Steinart PM. Microtubule disruption in keratinocytes induces cell-cell adhesion through activation of endogenous E-cadherin. *Mol Biol Cell*. 2001 Jul; 12(7): 1983-93.
 32. Kim S-Y, Bae C-D. Calpain inhibitors reduce the cornified cell envelope formation by inhibiting proteolytic processing of transglutaminase 1. *Exp Mol Med*. 1998 Dec; 30(4): 257-62.
 33. Kobayashi M, Yoshiki R, Sakabe J, Kabashima K, Nakamura M, Tokura Y.

- Expression of toll-like receptor 2, NOD2 and dectin-1 and stimulatory effects of their ligands and histamine in normal human keratinocytes. *Br J Dermatol.* 2009 Feb;160(2):297-304.
34. Kowanetz M, Valcourt U, Bergstrom R, Heldin C-H, Moustakas A. Id2 and Id3 define the potency of cell proliferation and differentiation responses to transforming growth factor β and bone morphogenetic protein. *Mol Cell Biol.* 2004 May; 24(10): 4241-54.
 35. Kurschat P, Bielenberg D, Rossignol-Tallandier M, Stahl A, Klagsbrun M. Neuron restrictive silencer factor NRSF/REST is a transcriptional repressor of neuropilin-1 and diminishes the ability of semaphorin 3A to inhibit keratinocyte migration. *J Biol Chem.* 2006; 281(5): 2721-9.
 36. Laux I, Jain A, Singh S, Agus DB. Epidermal growth factor receptor dimerization status determines skin toxicity to HER-kinase targeted therapies. *Br J Canc.* 2006; 94: 85-92.
 37. Lee B-K, Diebel E, Neukam FW, Wiltfang J, Ries J. Diagnostic and prognostic relevance of expression of human telomerase subunits in oral cancer. *Int J Oncol.* 2001; 19: 1063-8.
 38. Lee D-S, Quan G, Choi J-Y, Kim S-Y, Lee S-C. Chronic ultraviolet radiation modulates epidermal differentiation as it up-regulates transglutaminase 1 and its substrates. *Photodermatol Photoimmunol Photomed.* 2005 Feb; 21(1): 45-52.
 39. Lee ER, Kim JH, Kang YJ, Cho SG. The anti-apoptotic and anti-oxidant effect of eriodictyol on UV-induced apoptosis in keratinocytes. *Biol Pharm Bull.* 2007; 30(1):32-7.
 40. Lo RS, Witte ON. Transforming growth factor-beta activation promotes genetic context-dependent invasion of immortalized melanocytes. *Cancer Res.* 2008 Jun 1;68(11):4248-57.
 41. Loryman C, Mansbridge J. Inhibition of keratinocyte migration by lipopolysaccharide. *Wound Repair Regen.* 2008 Jan-Feb;16(1):45-51.
 42. McMillan JR, Akiyama M, Tanaka M, Yamamoto S, Goto M, Abe R, Sawamura D, Shimomura M, Shimizu H. Small-diameter porous poly (epsilon-caprolactone) films enhance adhesion and growth of human cultured epidermal keratinocyte and dermal fibroblast cells. *Tissue Eng.* 2007 Apr;13(4):789-98.
 43. Mildner M, Eckhart L, Lengauer B, Tschachler E. Hepatocyte growth factor/scatter factor inhibits UVB-induced apoptosis of human keratinocytes but not of keratinocyte-derived cell lines via the phosphatidylinositol 3-kinase/AKT pathway. *J Biol Chem.* 2002 Apr 19; 277(16): 14146-52.
 44. Mongan NP, Martin KM, Gudas LJ. The putative human stem cell marker, Rex-1 (Zfp42): structural classification and expression in normal human epithelial and carcinoma cell cultures. *Mol Carcinog.* 2006; 45(12):887-900.
 45. Müller-Schiffmann A, Beckmann J, Steger G. The E6 protein of the cutaneous human papillomavirus type 8 can stimulate the viral early and late promoters by distinct mechanisms. *J Virol.* 2006; 80(17):8718-28.
 46. Nakanishi G, Kim YS, Nakajima T, Jetten AM. Regulatory role for Kruppel-like zinc-finger protein Gli-similar 1 (Glis 1) in PMA-treated and psoriatic epidermis. *J Invest Dermatol.* 2006 Jan; 126(1): 49-60.
 47. Nelson AM, Gilliland KL, Cong Z, Thiboutot DM. 13-cis Retinoic acid induces apoptosis and cell cycle arrest in human SEB-1 sebocytes. *J Invest Dermatol.* 2006; 126(10):2178-89.
 48. Nicola AV, Hou J, Major EO, Straus SE. Herpes simplex virus type 1 enters human epidermal keratinocytes, but not neurons, via a pH-dependent endocytic pathway. *J Virol.* 2005 Jun; 79(12): 7609-16.
 49. Omata Y, Lewis JB, Rotenberg S, Lockwood PE, Messer RL, Noda M, Hsu SD, Sano H, Wataha JC. Intra- and extracellular reactive oxygen species generated by blue light. *J Biomed Mater Res A.* 2006; 77(3): 470-7.
 50. O'Neill AT, Monteiro-Riviere NA, Walker GM. Microfabricated curtains for controlled cell seeding in high throughput microfluidic systems. *Lab Chip.* 2009 Jun 21;9(12):1756-62.
 51. Oshikiri T, Miyamoto M, Hiraoka K, Shichinohe T, Kawarada Y, Kato K, Suzuoki M, Nakakubo Y, Kondo S, Dosaka-Akita H, Kasahara N, Katoh H. Transcriptional targeting of adenovirus vectors with the squamous cell carcinoma-specific antigen-2 promoter for selective apoptosis induction in lung cancer. *Cancer Gene Ther.* 2006;13(9):856-63.
 52. Ouyang W, Li J, Zhang D, Jiang BH, Huang DC. PI-3K/Akt signal pathway plays a crucial role in arsenite-induced cell proliferation of human keratinocytes through induction of

- cyclin D1. *J Cell Biochem.* 2007 Jul 1;101(4):969-78.
53. Papadimou E, Pavlidou D, Seraphin B, Tsambaos D, Drinas D. Retinoids inhibit human epidermal Keratinocyte Rnase P activity. *Biol Chem.* 2003 Mar; 384(3): 457-62.
 54. Perez DS, Handa RJ, Yang RS, Campaign JA. Gene expression changes associated with altered growth and differentiation in benzo[a]pyrene or arsenic exposed normal human epidermal keratinocytes. *J Appl Toxicol.* 2008 May;28(4):491-508.
 55. Petit-Frère C, Capulas E, Lyon DA, Norbury CJ, Lowe JE, Clingen pH, Riballo E, Green MHL, Arlett CF. Apoptosis and cytokine release induced by ionizing or ultraviolet B radiation in primary and immortalized human keratinocytes. *Carcinogen.* 2000; 21(6): 1087-95.
 56. Pivarsci A, Gombert M, Dieu-Nosjean M-C, Lauerma A, Kubitza R, Meller S, Ricker J, Muller A, Da Cunha L, Haahtela A, Sonkoly E, Fridman W-H, Alenius H, Kemeny L, Ruzicka T, Zlotnik A, Homey B. CC chemokine Ligand 18, an atopic dermatitis-associated and dendritic cell-derived chemokine, is regulated by staphylococcal products and allergen exposure. *J Immuno.* 2005; 173: 5810-7.
 57. Ridgway JBB, Ng E, Kern JA, Lee J, Brush J, Goddard A, Carter P. Identification of a human anti-CD55 single-chain Fv by subtractive panning of a phage library using tumor and nontumor cell lines. *Canc Res.* 1999 Jun 1; 59: 2718-23.
 58. Roomi MW, Monterrey JC, Kalinovsky T, Rath M, Niedzwiecki A. Distinct patterns of matrix metalloproteinase-2 and -9 expression in normal human cell lines. *Oncol Rep.* 2009 Mar; 21(3):821-6.
 59. Rouse JG, Haslauer CM, Loba EG, Monteiro-Riviere NA. Cyclic tensile strain increases interactions between human epidermal keratinocytes and quantum dot nanoparticles. *Toxicol In Vitro.* 2008 Mar;22(2):491-7.
 60. Sakai K, Akiyama M, Sugiyama-Nakagiri Y, McMillan JR, Sawamura D, Shimizu H. Localization of ABCA12 from Golgi apparatus to lamellar granules in human upper epidermal keratinocytes. *Exp Dermatol.* 2007 Nov;16(11):920-6.
 61. Sakurai N, Kuroiwa T, Ikeuchi H, Hiramatsu N, Maeshima A, Kaneko Y, Hiromura K, Nojima Y. Expression of IL-19 and its receptors in RA: potential role for synovial hyperplasia formation. *Rheumatology (Oxford).* 2008 Jun;47(6):815-20.
 62. Schäfer C, Borm B, Born S, Möhl C, Eibl EM, Hoffmann B. One step ahead: role of filopodia in adhesion formation during cell migration of keratinocytes. *Exp Cell Res.* 2009 Apr 15;315(7):1212-24.
 63. Schuster KM, Martens M, Goldenberg M, Tai C, Strande L, Hewitt CW. Production of a novel fibroblast-populated platelet matrix cocultured with keratinocytes. *Tissue Eng.* 2007 Jul;13(7):1493-500.
 64. Selby JC, Shannon MA. Apparatus for measuring the finite load-deformation behavior of a sheet of epithelial cells cultured on a mesoscopic freestanding elastomer membrane. *Rev Sci Instrum.* 2007 Sep;78(9):094301.
 65. Shen SI, Kotamraj PR, Bhattacharya S, Li X, Jasti BR. Synthesis and characterization of RGD-fatty acid amphiphilic micelles as targeted delivery carriers for anticancer agents. *J Drug Target.* 2007; 15(1):51-8.
 66. Shvedova AA, Tyurina YY, Tyurina VA, Kikuchi Y, Kagan VE, Quinn PJ. Quantitative analysis of phospholipids peroxidation and antioxidant protection in live human epidermal keratinocytes. *Biosci Repor.* 2001 Feb; 21(1): 33-43.
 67. Steiling H, Longet K, Moodycliffe A, Mansourian R, Bertschy E, Smola H, Mauch C, Williamson G. Sodium-dependent vitamin C transporter isoforms in skin: Distribution, kinetics, and effect of UVB-induced oxidative stress. *Free Radic Biol Med.* 2007 Sep 1;43(5):752-62.
 68. Steinert PM, Marekov LN. Initiation of assembly of the cell envelope barrier structure of stratified squamous epithelia. *Am Soc Cell Biol.* 1999; 10: 4247-61.
 69. Stoff A, Rivera AA, Banerjee NS, Mathis JM, Espinosa-de-los-Monteros A, Le LP, De la Torre JI, Vasconez LO, Broker TR, Richter DF, Stoff-Khalili MA, Curiel DT. Strategies to enhance transductional efficiency of adenoviral-based gene transfer to primary human fibroblasts and keratinocytes as a platform in dermal wounds. *Wound Repair Regen.* 2006;14(5):608-17.
 70. Subbarayan V, Xu X-C, Kim J, Yang P, Hoque A, Sabichi AL, Llansa N, Mendoza G, Logothetis CJ, Newman RA, Lippman SM, Menter DG. Inverse relationship between 15-lipoxygenase-2 and PPAR- γ gene expression in normal epithelia compared with tumor epithelia. *Neoplas.* 2005 Mar; 7(3): 280-93.

71. Tani K, Adachi M, Nakamura Y, Kano R, Makimura K, Hasegawa A, Kanda N, Watanabe S. The effect of dermatophytes on cytokine production by human keratinocytes. *Arch Dermatol Res.* 2007 Oct;299(8):381-7.
72. Tekos A, Prodromaki E, Papadimou E, Pavlidou D, Tsambaos D, Drainas D. Aminoglycosides suppress tRNA processing in human epidermal keratinocytes in vitro. *Skin Pharmacol Appl Skin Physiol.* 2003; 16(4): 252-8.
73. Tominaga M, Ogawa H, Takamori K. Possible roles of epidermal opioid systems in pruritus of atopic dermatitis. *J Invest Dermatol.* 2007 Sep;127(9):2228-35.
74. Tran NQ, Crowe DL. Regulation of the human involucrin gene promoter by co-activator proteins. *Biochem J.* 2004; 381: 267-73.
75. Trouba KJ, Geisenhoffer KM, Germolec DR. Sodium arsenite-induced stress-related gene expression in normal human epidermal, HaCaT, and HEL-30 keratinocytes. *Env Health Perspect.* 2002 oct; 110(5): 761-6.
76. Wang YH, Angkasekwinai P, Lu N, Voo KS, Arima K, Hanabuchi S, Hippe A, Corrigan CJ, Dong C, Homey B, Yao Z, Ying S, Huston DP, Liu YJ. IL-25 augments type 2 immune responses by enhancing the expansion and functions of TSLP-DC-activated Th2 memory cells. *J Exp Med.* 2007 Aug 6;204(8):1837-47.
77. Wataha JC, Lewis JB, Lockwood PE, Hsu S, Messer RL, Rueggeberg FA, Bouillaguet S. Blue light differentially modulates cell survival and growth. *J Dent Res.* 2004; 83(2): 104-8.
78. Watanabe S, Kano R, Sato H, Nakamura Y, Hasegawa A. The effects of Malassezia yeasts on cytokine production by human keratinocytes. *J Invest Dermatol.* 2001; 116: 769-73.
79. Wu F, Burns FJ, Zhang R, Uddin AN, Rossman TG. Arsenite-induced alterations of DNA photodamage repair and apoptosis after solar-simulation UVR in mouse keratinocytes in vitro. *Env Health Perspect.* 2005; 113: 983-6.
80. Yamamoto T, Lewis J, Wataha J, Dickinson D, Singh B, Bollag WB, Ueta E, Osaki T, Athar M, Schuster G, Hsu S. Roles of catalase and hydrogen peroxide in green tea polyphenol-induced chemopreventive effects. *J Pharmacol Exp Therap.* 2004; 308: 317-23.
81. Yang Z-Y, Huang Y, Ganesh L, Leung K, Kong W-P, Schqartz O, Subbarao K, Nabel GJ. pH-dependent entry of severe acute respiratory syndrome coronavirus is mediated by the spike glycoprotein and enhanced by dendritic cell transfer through DC-SIGN. *J Virol.* 2004 Jun; 78(11): 5642-50.
82. Yarosh DB, Galvin JW, Nay SL, Peña AV, Canning MT, Brown DA. Anti-inflammatory activity in skin by biomimetic of *Evodia rutaecarpa* extract from traditional Chinese medicine. *J Dermatol Sci.* 2006; 42(1):13-21.
83. Yasukawa K, Sawamura D, Goto M, Nakamura H, Shimizu H. Histone deacetylase inhibitors preferentially augment transient transgene expression in human dermal fibroblasts. *Br J Dermatol.* 2007 Oct;157(4):662-9.
84. Yin W, Cheepala S, Roberts JN, Syson-Chan K, DiGiovanni J, Clifford JL. Active Stat3 is required for survival of human squamous cell carcinoma cells in serum-free conditions. *Mol Cancer.* 2006;5:15.
85. Yoon K, Smart RC. C/EBPalpha is a DNA damage-inducible p53-regulated mediator of the G1 checkpoint in Keratinocytes. *Mol Cell Biol.* 2004 Dec; 24(24): 10650-60.
86. Yoshida A, Kanno H, Watabe D, Akasaka T, Sawai T. The role of heparin-binding EGF-like growth factor and amphiregulin in the epidermal proliferation of psoriasis in cooperation with TNFalpha. *Arch Dermatol Res.* 2008 Jan;300(1):37-45.
87. Zhang LW, Yu WW, Colvin VL, Monteiro-Riviere NA. Biological interactions of quantum dot nanoparticles in skin and in human epidermal keratinocytes. *Toxicol Appl Pharmacol.* 2008 Apr 15;228(2):200-11.
88. Zucchini C, Strippoli P, Biolchi A, Solmi R, Lenzi L, D'Addabbo P, Carinci P, Valvassori. The human TruB family of pseudouridine synthase genes, including the Dyskeratosis Congenita 1 gene and the novel member TRUB1. *Int J Mol Med.* 2003; 11: 697-704.

Media

1. Ansari KM, Rundhaug JE, Fischer SM. Multiple signaling pathways are responsible for prostaglandin E2-induced murine keratinocyte proliferation. *Mol Cancer Res.* 2008 Jun;6(6):1003-16.
2. Bajpai M, Liu J, Geng X, Souza RF, Amenta PS, Das KM. Repeated exposure to acid

- and bile selectively induces colonic phenotype expression in a heterogeneous Barrett's epithelial cell line. *Lab Invest.* 2008 Jun;88(6):643-51.
3. Baten A, Sakamoto K, Shamsuddin AM. Long-term culture of normal human colonic epithelial cells in vitro. *FASEB Journal.* 1992; 6: 2726-34.
 4. Calkins CC, Setzer SV, Jennings JM, Summers S, Tsunoda K, Amagai M, Kowalczyk AP. Desmoglein endocytosis and desmosome disassembly are coordinated responses to pemphigus autoantibodies. *J Biol Chem.* 2006; 281(11):7623-34.
 5. Castilho RM, Squarize CH, Patel V, Millar SE, Zheng Y, Molinolo A, Gutkind JS. Requirement of Rac1 distinguishes follicular from interfollicular epithelial stem cells. *Oncogene.* 2007 Aug 2;26(35):5078-85.
 6. Chamulitrat W, Huber A, Riedel HD, Stremmel W. Nox1 induces differentiation resistance in immortalized human keratinocytes generating cells that express simple epithelial keratins. *J Invest Dermatol.* 2007 Sep;127(9):2171-83.
 7. Culp TD, Budgeon LR, Marinkovich MP, Meneguzzi G, Christensen ND. Keratinocyte-secreted laminin 5 can function as a transient receptor for human papillomaviruses by binding virions and transferring them to adjacent cells. *J Virol.* 2006; 80(18):8940-50.
 8. Feagins LA, Zhang HY, Hormi-Carver K, Quinones MH, Thomas D, Zhang X, Terada LS, Spechler SJ, Ramirez RD, Souza RF. Acid has antiproliferative effects in nonneoplastic Barrett's epithelial cells. *Am J Gastroenterol.* 2007 Jan;102(1):10-20.
 9. Ishibashi Y, Sugita T, Nishikawa A. Cytokine secretion profile of human keratinocytes exposed to *Malassezia* yeasts. *FEMS Immunol Med Microbiol.* 2006; 48(3): 400-9.
 10. Jaiswal KR, Morales CP, Feagins LA, Gandia KG, Zhang X, Zhang HY, Hormi-Carver K, Shen Y, Elder F, Ramirez RD, Sarosi GA Jr, Spechler SJ, Souza RF. Characterization of telomerase-immortalized, non-neoplastic, human Barrett's cell line (BAR-T). *Dis Esophagus.* 2007;20(3):256-64.
 11. Kim HJ, Abdelkader N, Katz M, McLane JA. 1,25-dihydroxy-vitamin-D3 enhances antiproliferative effect and transcription of TGF-beta1 on human Keratinocytes in culture. *J Cell Physiol.* 1992 Jun; 151(3): 579-87.
 12. Kim TI, Chung JL, Hong JP, Min K, Seo KY, Kim EK. Bevacizumab application delays epithelial healing in rabbit cornea. *Invest Ophthalmol Vis Sci.* 2009 May 20. [Epub ahead of print]
 13. Kim YM, Cao D, Reed W, Wu W, Jaspers I, Tal T, Bromberg PA, Samet JM. Zn2+-induced NF-kappaB-dependent transcriptional activity involves site-specific p65/RelA phosphorylation. *Cell Signal.* 2007 Mar;19(3):538-46.
 14. Kostan J, Gregor M, Walko G, Wiche G. Plectin Isoform-dependent Regulation of Keratin-Integrin {alpha}6{beta}4 Anchorage via Ca2+/Calmodulin. *J Biol Chem.* 2009 Jul 3;284(27):18525-36.
 15. Lee C, Wooldridge TR, Laimins LA. Analysis of the roles of E6 binding to E6TP1 and nuclear localization in the human papillomavirus type 31 life cycle. *Virology.* 2007 Feb 5;358(1):201-10.
 16. Li J, Raghunath M, Tan D, Lareu RR, Chen Z, Beuerman RW. Defensins HNP1 and HBD2 stimulation of wound-associated responses in human conjunctival fibroblasts. *Invest Ophthalmol Vis Sci.* 2006; 47(9):3811-9.
 17. Luo D, Lin XF, Min W, Ma QH, Gu N, Jin SL, Wang DG. Photoprotection by tocopherol submicron emulsion against UV-mediated damage in HaCaT cells. *Methods Find Exp Clin Pharmacol.* 2007 Apr;29(3):185-9.
 18. Marconi A, Dallaglio K, Lotti R, Vaschieri C, Truzzi F, Fantini F, Pincelli C. Survivin identifies keratinocyte stem cells and is downregulated by anti-beta1 integrin during anoikis. *Stem Cells.* 2007; 25(1):149-55.
 19. Marinari B, Ballaro C, Koster MI, Giustizieri ML, Moretti F, Crosti F, Papoutsaki M, Karin M, Alema S, Chimenti S, Roop DR, Costanzo A. IKKalpha is a p63 transcriptional target involved in the pathogenesis of ectodermal dysplasias. *J Invest Dermatol.* 2009 Jan;129(1):60-9.
 20. Marinkovich MP, Lunstrum GP, Keene DR, Burgeson RE. The dermal-epidermal junction of human skin contains a novel laminin variant. *J Cell Biol.* 1992; 119(3): 695-703.
 21. Milligan SG, Veerapraditsin T, Ahamet B, Mole S, Graham SV. Analysis of novel human papillomavirus type 16 late mRNAs

- in differentiated W12 cervical epithelial cells. *Virology*. 2007 Mar 30;360(1):172-81.
22. Naim R, Sadick H, Bayerl C, Riedel F, Schafer C, Bran G, Hormann K. Heptocyte growth factor/scatter factor induces VEGF in human external auditory canal cholesteatoma cell culture. *Int J Mol Med*. 2005; 15: 67-71.
 23. Pillai S, Bikle DD, Hincenbergs M, Elias PM. Biochemical and morphological characterization of growth and differentiation of normal human neonatal keratinocytes in a serum-free medium. *J Cell Physiol*. 1988; 134: 229-37.
 24. Reilly CA, Johansen ME, Lanza DL, Lee L, Lim J-O, Yost GS. Calcium-dependent and independent mechanisms of capsaicin receptor (TRPV1)-mediated cytokine production and cell death in human bronchial epithelial cells. *J Biochem Mol Toxicol*. 2005; 19(4): 266-75.
 25. Resau JH, Phelps PC, Zhu SM, Smoot D, Lee HK, Cottrell JR, Hudson EA, Elliget KA. Long-term culture of human esophageal explants and cells. *Cytotechnol*. 1990 Jan; 3(1): 61-73.
 26. Rice RH, Steinmann KE, deGraffenried L, Qin Q, Taylor N, Schlegel R. Elevation of cell cycle control proteins during spontaneous immortalization of human keratinocytes. *Mol Biol Cell*. 1993; 4: 185-94.
 27. Robertson DM, Ho SI, Cavanagh HD. Characterization of DeltaNp63 isoforms in normal cornea and telomerase-immortalized human corneal epithelial cells. *Exp Eye Res*. 2008 Apr;86(4):576-85.
 28. Robertson DM, Li L, Fisher S, Pearce VP, Shay JW, Wright WE, Cavanagh HD, Jester JV. Characterization of growth and differentiation in a telomerase-immortalized human corneal epithelial cell line. *Invest Ophthalmol Vis Sci*. 2005; 46: 470-8.
 29. Sharma GD, Kakazu A, Bazan HE. Protein kinase C alpha and epsilon differentially modulate hepatocyte growth factor-induced epithelial proliferation and migration. *Exp Eye Res*. 2007 Aug;85(2):289-97.
 30. Shin KH, Kim RH, Kang MK, Kim RH, Kim SG, Lim PK, Yochim JM, Baluda MA, Park NH. p53 promotes the fidelity of DNA end-joining activity by, in part, enhancing the expression of heterogeneous nuclear ribonucleoprotein G. *DNA Repair (Amst)*. 2007 Jun 1;6(6):830-40.
 31. Tiala I, Wakkinen J, Suomela S, Puolakkainen P, Tammi R, Forsberg S, Rollman O, Kainu K, Rozell B, Kere J, Saarialho-Kere U, Elomaa O. The PSORS1 locus gene CCHCR1 affects keratinocyte proliferation in transgenic mice. *Hum Mol Genet*. 2008 Apr 1;17(7):1043-51.
 32. Varani J, Bhagavathula N, Aslam MN, Fay K, Warner RL, Hanosh A, Barron AG, Miller RA. Inhibition of retinoic acid-induced skin irritation in calorie-restricted mice. *Arch Dermatol Res*. 2008 Jan;300(1):27-35.
 33. Varani J, Fay K, Perone P. MDI 301, a non-irritating retinoid, induces changes in human skin that underlie repair. *Arch Dermatol Res*. 2007; 298(9):439-48.
 34. Veranth JM, Moss TA, Chow JC, Labban R, Nichols WK, Walton JC, Watson JG, Yost GS. Correlation of in vitro cytokine responses with the chemical composition of soil-derived particulate matter. *Env Health Perspec*. 2006 Mar; 114(3): 341-9.
 35. Vigneswaran N, Baucum DC, Wu J, Lou Y, Bouquot J, Muller S, Zacharias W. Repression of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) but not its receptors during oral cancer progression. *BMC Cancer*. 2007 Jun 25;7:108.
 36. Williams R, Zou X, Hoyle GW. Tachykinin-1 receptor stimulates proinflammatory gene expression in lung epithelial cells through activation of NF-kappaB via a G(q)-dependent pathway. *Am J Physiol Lung Cell Mol Physiol*. 2007 Feb;292(2):L430-7.
 37. Yi X-J, Wang Y, Yu F-S. Corneal epithelial tight junctions and their response to lipopolysaccharide challenge. *Invest Ophthalmol Vis Sci*. 2000; 41: 4093-400.
 38. Yilmaz O, Verbeke P, Lamont RJ, Ojcius DM. Intercellular spreading of *Porphyromonas gingivalis* infection in primary gingival epithelial cells. *Infect Immun*. 2006;74(1):703-10.
 39. Yoo H, Kim SJ, Kim Y, Lee H, Kim TY. Insulin-like growth factor-II regulates the 12-lipoxygenase gene expression and promotes cell proliferation in human keratinocytes via the extracellular regulatory kinase and phosphatidylinositol 3-kinase pathways. *Int J Biochem Cell Biol*. 2007;39(6):1248-59.

* References not specifically citing the use of Lonza cells, media, or reagents in their research.

+ Denotes sections containing only the articles published within the last ten years.