

Product References

Clonetics® Renal Cells

Cells

RPTEC⁺

1. Abend JR, Low JA, Imperiale MJ. Inhibitory effect of gamma interferon on BK virus gene expression and replication. *J Virol.* 2007 Jan;81(1):272-9.
2. Ami Y, Shimazui T, Akaza H, Uematsu N, Yano Y, Tsujimoto G, Uchida K. Gene expression profiles correlate with the morphology and metastasis characteristics of renal cell carcinoma cells. *Oncol Rep.* 2005; 13: 75-80.
3. Asakuma J, Sumitomo M, Asano T, Asano T, Hayakawa M. Selective Akt inactivation and tumor necrosis factor-related apoptosis-inducing ligand sensitization of renal cancer cells by low concentrations of paclitaxel. *Canc Res.* 2003 Mar 15; 63: 1365-70.
4. Beqaj S, Shah AM, Ryan JM. Identification of cells responsible for urokinase-type plasminogen activator synthesis and secretion in human diploid kidney cell cultures. *In Vitro Cell Dev Biol Anim.* 2004 Mar-Apr;40(3-4):102-7.
5. Bernhoff E, Gutteberg TJ, Sandvik K, Hirsch HH, Rinaldo CH. Cidofovir inhibits polyomavirus BK replication in human renal tubular cells downstream of viral early gene expression. *Am J Transplant.* 2008 Jul;8(7):1413-22.
6. Braun MC, Kelly CN, Prada AE, Mishra J, Chand D, Devarajan P, Zahedi K. Human PRMT5 expression is enhanced during in vitro tubule formation and after in vivo ischemic injury in renal epithelial cells. *Am J Nephrol.* 2004; 24(2): 250-7.
7. Campbell CE, Kuriyan NP, Tackley RR, Caulfield MJ, Tubbs R, Finke J, Williams BRG. Constitutive expression of wilms tumor suppressor gene(WT1) in renal cell carcinoma. *Int J Canc.* 1999; 78(2): 182-8.
8. Chan T-M, Leung JK-H, Ho S K-N, Yung S. Mesangial cell-binding anti-DNA antibodies in patients with systemic lupus erythematosus. *J Am Soc Nephrol.* 2002; 13: 1219-29.
9. Chen Q, Shinohara N, Abe T, Watanabe T, Nonomura K, Koyanagi T. Significance of COX-2 expression in human renal cell carcinoma cell lines. *Int J Canc.* 2004; 108(6): 825-32.
10. Chi JT, Wang Z, Nuyten DS, Rodriguez EH, Schaner ME, Salim A, Wang Y, Kristensen GB, Helland A, Børresen-Dale AL, Giaccia A, Longaker MT, Hastie T, Yang GP, van de Vijver MJ, Brown PO. Gene expression programs in response to hypoxia: cell type specificity and prognostic significance in human cancers. *PLoS Med.* 2006 Mar; 3(3):e47.
11. Cihlar T, Ray AS, Bojamra CG, Zhang L, Hui H, Laflamme G, Vela JE, Grant D, Chen J, Myrick F, White KL, Gao Y, Lin KY, Douglas JL, Parkin NT, Carey A, Pakdaman R, Mackman RL. Design and profiling of GS-9148, a novel nucleotide analog active against nucleoside-resistant variants of human immunodeficiency virus type 1, and its orally bioavailable phosphonoamidate prodrug, GS-9131. *Antimicrob Agents Chemother.* 2008 Feb;52(2):655-65.
12. Datta D, Dormond O, Basu A, Briscoe DM, Pal S. Heme oxygenase-1 modulates the expression of the anti-angiogenic chemokine CXCL-10 in renal tubular epithelial cells. *Am J Physiol Renal Physiol.* 2007 Oct;293(4):F1222-30.
13. de Paulsen N, Brychzy A, Fournier M-C, Klausner RD, Gnarra JR, Pause A, Lee S. Role of transforming growth factor- α in von Hippel-Lindau (VHL)^{-/-} clear cell renal

- carcinoma cell proliferation: a possible mechanism coupling VHL tumor suppressor inactivation and tumorigenesis. PNAS. 2001 Feb 13; 98(4): 1387-92.
14. Drumm K, Kress TR, Gassner B, Krug AW, Gekle M. Aldosterone stimulates activity and surface expression of NHE3 in human primary renal proximal tubule epithelial cells (RPTEC). *Cell Physiol Biochem*. 2006; 17: 21-8.
 15. Dudas PL, Argentieri RL, Farrell FX. BMP-7 fails to attenuate TGF-beta1-induced epithelial-to-mesenchymal transition in human proximal tubule epithelial cells. *Nephrol Dial Transplant*. 2009 May;24(5):1406-16.
 16. Elgadi KM, Meguid RA, Qian M, Souba WW, Abcouwer SF. Cloning and analysis of unique human glutaminase isoforms generated by tissue-specific alternative splicing. *Physiol Genomics*. 1999 Aug 31; 1(2): 51-62.
 17. Fujii J, Wood K, Matsuda F, Carneiro-Filho BA, Schlegel KH, Yutsudo T, Binnington-Boyd B, Lingwood CA, Obata F, Kim KS, Yoshida S, Obrig T. Shiga toxin 2 causes apoptosis in human brain microvascular endothelial cells via C/EBP homologous protein. *Infect Immun*. 2008 Aug;76(8):3679-89.
 18. Gerstung M, Roth T, Dienes HP, Licht C, Fries JW. Endothelin-1 induces NF-kappaB via two independent pathways in human renal tubular epithelial cells. *Am J Nephrol*. 2007;27(3):294-300.
 19. Girard C, Barbier O, Beilleux G, El-Alfy M, Bélanger A. Human uridine diphosphate-glucuronosyltransferase UGT2B7 conjugates mineralocorticoid and glucocorticoid metabolites. *Endocrinol*. 2003; 144: 2659-68.
 20. Gosert R, Rinaldo CH, Funk GA, Egli A, Ramos E, Drachenberg CB, Hirsch HH. Polyomavirus BK with rearranged noncoding control region emerge in vivo in renal transplant patients and increase viral replication and cytopathology. *J Exp Med*. 2008 Apr 14;205(4):841-52.
 21. Hergovich A, Lisztwan J, Thoma CR, Wirbelauer C, Barry RE, Krek W. Priming-dependent phosphorylation and regulation of the tumor suppressor pVHL by glycogen synthase kinase 3. *Mol Cell Biol*. 2006; 26(15): 5784-96.
 22. Hughes AK, Schmid DI, Kohan DE. Sex steroids do not affect shigatoxin cytotoxicity on human renal tubular or glomerular cells. *BMC Nephrol*. 2002; 3: 6-13.
 23. Hughes AK, Stricklett PK, Kohan DE. Cytotoxic effect of Shiga toxin-1 on human proximal tubule cells. *Kidney Int*. 1998 Aug;54(2):426-37.
 24. Juang SH, Lung CC, Hsu PC, Hsu KS, Li YC, Hong PC, Shiah HS, Kuo CC, Huang CW, Wang YC, Huang L, Chen TS, Chen SF, Fu KC, Hsu CL, Lin MJ, Chang CJ, Ashendel CL, Chan TC, Chou KM, Chang JY. D-501036, a novel selenophene-based triheterocycle derivative, exhibits potent in vitro and in vivo antitumoral activity which involves DNA damage and ataxia telangiectasia-mutated nuclear protein kinase activation. *Mol Cancer Ther*. 2007; 6(1):193-202.
 25. Junaid A, Amara FM. Osteopontin: correlation with interstitial fibrosis in human diabetic kidney and PI3-kinase-mediated enhancement of expression by glucose in human proximal tubular epithelial cells. *Histopathol*. 2004 Feb; 44(2): 136-46.
 26. Kamal A, Thao L, Sensintaffar J, Zhang L, Boehm MF, Fritz LC, Burrows FJ. A high-affinity conformation of Hsp90 confers tumour selectivity on Hsp90 inhibitors. *Nature*. 2003 Sep 25; 425: 407-10.
 27. Keshi H, Sakamoto T, Kawai T, Ohtani K, Katoh T, Jang SJ, Motomura W, Yoshizaki T, Fukuda M, Koyama S, Fukuzawa J, Fukuoh A, Yoshida I, Suzuki Y, Wakamiya N. Identification and characterization of a novel human collectin CL-K1. *Microbiol Immunol*. 2006;50(12):1001-13.
 28. Königs M, Lenczyk M, Schwerdt G, Holzinger H, Gekle M, Humpf HU. Cytotoxicity, metabolism and cellular uptake of the mycotoxin deoxynivalenol in human proximal tubule cells and lung fibroblasts in primary culture. *Toxicology*. 2007 Oct 30;240(1-2):48-59.
 29. Königs M, Mulac D, Schwerdt G, Gekle M, Humpf HU. Metabolism and cytotoxic effects of T-2 toxin and its metabolites on human cells in primary culture. *Toxicology*. 2009 Apr 28;258(2-3):106-15.
 30. Laping NJ, Grygielko E, Mathur A, Butter S, Bomberger J, Tweed C, Martin W, Fornwald J, Lehr R, Harling J, Gaster L, Callahan JF, Olson BA. Inhibition of transforming growth factor (TGF)- β 1-induced extracellular matrix

- with a novel inhibitor of the TGF- β type I receptor kinase activity: SB-431542. *Mol Pharmacol.* 2002; 62: 58-64.
31. Laxmanan S, Datta D, Geehan C, Briscoe DM, Pal S. CD40: a mediator of pro- and anti- inflammatory signals in renal tubular epithelial cells. *J Am Soc Nephrol.* 2005; 16(9): 2714-23.
 32. Leuenberger D, Andresen PA, Gosert R, Binggeli S, Ström EH, Bodaghi S, Rinaldo CH, Hirsch HH. Human polyomavirus type 1 (BK virus) agnoprotein is abundantly expressed but immunologically ignored. *Clin Vaccine Immunol.* 2007 Aug;14(8):959-68.
 33. Ma Q, Devarajan P. Induction of proapoptotic Daxx following ischemic acute kidney injury. *Kidney Int.* 2008 Aug;74(3):310-8.
 34. Mahon C, Liang B, Tikhanovich I, Abend JR, Imperiale MJ, Nasheuer HP, Folk WR. Restriction of human polyomavirus BK virus DNA replication in murine cells and extracts. *J Virol.* 2009 Jun;83(11):5708-17.
 35. Marini P, Schmid A, Jendrossek V, Faltin H, Daniel PT, Budach W, Belka C. Irradiation specifically sensitizes solid tumour cell lines to TRAIL mediated apoptosis. *BMC Canc.* 2005; 5(5).
 36. Moriyama T, Marquez JP, Wakatsuki T, Sorokin A. Caveolar endocytosis is critical for BK virus infection of human renal proximal tubular epithelial cells. *J Virol.* 2007 Aug;81(16):8552-62.
 37. Moriyama T, Sorokin A. Intracellular trafficking pathway of BK Virus in human renal proximal tubular epithelial cells. *Virology.* 2008 Feb 20;371(2):336-49.
 38. Nakagawa N, Yao M, Baba M, Kato S, Kishida T, Hattori K, Nagashima Y, Kubota Y. Inactivation of von Hippel-Lindau gene induces constitutive phosphorylation of MET protein in clear cell renal carcinoma. *Canc Res.* 2006; 66: 3699-705.
 39. Nguyen HT, Hsieh MH, Gaborro A, Tinloy B, Phillips C, Adam RM. JNK/SAPK and p38 SAPK-2 mediate mechanical stretch-induced apoptosis via caspase-3 and -9 in NRK-52E renal epithelial cells. *Nephron Exp Nephrol.* 2006; 102(2):e49-61.
 40. Nyström J, Hultenby K, Ek S, Sjölund J, Axelson H, Jirström K, Saleem MA, Nilsson K, Johansson ME. CRIM1 is localized to the podocyte filtration slit diaphragm of the adult human kidney. *Nephrol Dial Transplant.* 2009 Jul;24(7):2038-44.
 41. Ovcharenko D, Jarvis R, Hunicke-Smith S, Kelnar K, Brown D. High-throughput screening in vitro: from cell lines to primary cells. *RNA.* 2005; 11: 985-93.
 42. Park S-W, Bae J-S, Kim K-S, Park S-H, Lee B-H, Choi J-Y, Ha S-W, Kim Y-L, Kwon T-H, Kim I-S, Park R-W. Beta ig-h3 promotes renal proximal tubular epithelial cell adhesion, migration, and proliferation through the interaction with $\alpha 3\beta 1$ integrin. *Exp Mol Med.* 2004 Jun; 36(3): 211-9.
 43. Peruzzi B, Athauda G, Bottaro DP. The von Hippel-Lindau tumor suppressor gene product represses oncogenic beta-catenin signaling in renal carcinoma cells. *Proc Natl Acad Sci U S A.* 2006 Sep 26; 103(39): 14531-6.
 44. Prozialeck WC, Edwards JR, Lamar PC. Evaluation of the barrier properties of a commercially available renal epithelial cell system. *Toxicol In Vitro.* 2007 Apr;21(3):533.
 45. Psotka MA, Obata F, Kolling GL, Gross LK, Saleem MA, Satchell SC, Mathieson PW, Obrig TG. Shiga toxin 2 targets the murine renal collecting duct epithelium. *Infect Immun.* 2009 Mar;77(3):959-69.
 46. Rey O, Young SH, Yuan J, Slice L, Rozengurt E. Amino acid-stimulated Ca^{2+} oscillations produced by the Ca^{2+} -sensing receptor are mediated by a phospholipase C/inositol 1,4,5-triphosphate-independent pathway that requires G_{12} , rho, filament-A, and the actin cytoskeleton. *J Biol Chem.* 2005; 280(24): 22875-82.
 47. Salahudeen AK, Haider N, Jenkins J, Joshi M, Patel H, Huang H, Yang M, Zhe H. Antia apoptotic properties of erythropoiesis-stimulating proteins in models of cisplatin-induced acute kidney injury. *Am J Physiol Renal Physiol.* 2008 Jun;294(6):F1354-65.
 48. Schmid DI, Kohan DE. Effect of shigatoxin-1 on arachidonic acid release by human glomerular epithelial cells. *Kidney Int.* 2001 Sep;60(3):1026-36.
 49. Schwerdt G, Holzinger H, Sauvant C, Königs M, Humpf HU, Gekle M. Long-term effects of ochratoxin A on fibrosis and cell death in human proximal tubule or fibroblast cells in primary culture. *Toxicology.* 2007 Mar 22;232(1-2):57-67.
 50. Schwerdt G, Königs M, Holzinger H, Humpf HU, Gekle M. Effects of the mycotoxin

LONZA

- fumonisin B(1) on cell death in human kidney cells and human lung fibroblasts in primary culture. *J Appl Toxicol.* 2009 Mar;29(2):174-82.
51. Shikano N, Kawai K, Nakajima S, Nishii R, Flores II LG, Kubodera A, Kubota N, Ishikawa N, Saji H. Renal accumulation and excretion of radioiodinated 3-iodo- α -methyl-L-tyrosine. *Ann Nuc Med.* 2004; 18(3): 263-70.
52. Shimazui T, Ami Y, Yoshikawa K, Uchida K, Kojima T, Oikawa T, Nakamura K, Honda N, Hinotsu S, Miyazaki J, Kunita N, Akaza H. Prediction of in vitro response to interferon-alpha in renal cell carcinoma cell lines. *Cancer Sci.* 2007 Apr;98(4):529-34. Epub 2007 Jan 31.
53. Singh KA, Kampen RL, Hoffmann SC, Eldaif SM, Kirk AD. Renal epithelial cell-derived monocyte colony stimulating factor as a local informant of renal injury and means of monocyte activation. *Transpl Int.* 2009 Jul;22(7):730-7.
54. Singh R, Browning JL, Abi-Habib R, Wong K, Williams SA, Merchant R, Denmeade SR, Buckley TJ, Frankel AE. Recombinant prostate-specific antigen proaerolysin shows selective protease sensitivity and cell cytotoxicity. *Anticancer Drugs.* 2007 Aug;18(7):809-16.
55. Slattery C, Ryan MP, McMorrow T. Protein kinase C beta overexpression induces fibrotic effects in human proximal tubular epithelial cells. *Int J Biochem Cell Biol.* 2008;40(10):2218-29.
56. Son D, Kojima I, Inagi R, Matsumoto M, Fujita T, Nangaku M. Chronic hypoxia aggravates renal injury via suppression of Cu/Zn-SOD: a proteomic analysis. *Am J Physiol Renal Physiol.* 2008 Jan;294(1):F62-72.
57. Stec DE, Bishop C, Rimoldi JM, Poreddy SR, Vera T, Salahudeen AK. Carbon monoxide (CO) protects renal tubular epithelial cells against cold-rewarm apoptosis. *Ren Fail.* 2007;29(5):543-8.
58. Thoma CR, Frew IJ, Hoerner CR, Montani M, Moch H, Krek W. pVHL and GSK3beta are components of a primary cilium-maintenance signalling network. *Nat Cell Biol.* 2007 May;9(5):588-95.
59. Thomas-Tikhonenko A, Viard-Leveugle I, Dews M, Wehrli P, Sevignani C, Yu D, Ricci S, el-Deiry W, Aronow B, Kaya G, Saurat J-H, French LE. Myc-transformed epithelial cells down-regulate clusterin, which inhibits their growth in vitro and carcinogenesis in vivo. *Canc Res.* 2004 May 01; 64: 3126-36.
60. Vanoosten RL, Earel JK, Griffith TS. Enhancement of Ad5-TRAIL cytotoxicity against renal cell carcinoma with histone deacetylase inhibitors. *Canc Gene Ther.* 2006 Jun; 13(6): 628-32.
61. VanOosten RL, Earel JK, Griffith TS. Enhancement of Ad5-TRAIL cytotoxicity against renal cell carcinoma with histone deacetylase inhibitors. *Cancer Gene Ther.* 2006;13(6):628-32.
62. Vanoosten RL, Moore JM, Karacay B, Griffith TS. Histone deacetylase inhibitors modulate renal cell carcinoma sensitivity to TRAIL/Apo-2L-induced apoptosis by enhancing TRAIL-R2 expression. *Canc Biol Ther.* 2005; 4(10): 1104-12.
63. Wilson PD, Hovater JS, Casey CC, Fortenberry JA, Schwiebert EM. ATP release mechanisms in primary cultures of epithelia derived from the cysts of polycystic kidneys. *Am Soc of Nephrol.* 1999; 10.
64. Woods JS, Dieguez-Acuña, Ellis ME, Kushleika J, Simmonds PL. Attenuation of nuclear factor kappa B (NK- κ B) promotes apoptosis of kidney epithelial cells: a potential mechanism of mercury-induced nephrotoxicity. *Env Health Perspect.* 2002 Oct; 110(5): 819-22.
65. Yang Z-Y, Huang Y, Ganesh L, Leung K, Kong W-P, Schqartz O, Subbarao K, Nabel GJ. pH-dependent entry of sever acute respiratory syndrome coronavirus is mediated by the spike glycoprotein and enhanced by dendritic cell transfer through DC-SIGN. *J Virol.* 2004; 78(11): 5642-50.

HRCE

1. Beqaj S, Shah AM, Ryan JM. Identification of cells responsible for urokinase-type plasminogen activator synthesis and secretion in human diploid kidney cell cultures. *In Vitro Cell Dev Biol Anim.* 2004 Mar-Apr;40(3-4):102-7.
2. Black JB, Burns DA, Goldsmith CS, Feorino PM, Kite-Powell K, Schinazi RF, Krug PW, Pellett PE. Biologic properties of human herpesvirus 7 strain SB. *Virus Res.* 1997 Nov;52(1):25-41.
3. Braam B, Allen P, Benes E, Koomans HA, Navar LG, Hammond T. Human proximal tubular cell responses to angiotensin II

- analyzed using DNA microarray. *Eur J Pharmacol.* 2003 Mar 19;464(2-3):87-94.
4. Chen R-H, McCormick F. Selective targeting to the hyperactive β -catenin/T-cell factor pathway in colon cancer cells. *Canc Res.* 2001 Jun 1; 61: 4445-9.
 5. Hanaoka K, Guggino WB. cAMP regulates cell proliferation and cyst formation in autosomal polycystic kidney disease cells. *J Am Soc Nephrol.* 2000; 11: 1179-87.
 6. Itani OA, Liu KZ, Cornish KL, Campbell JR, Thomas CP. Glucocorticoids stimulate human sgk1 gene expression by activation of a GRE in its 5'-flanking region. *Am J Physiol Endocrinol Metab.* 2002; 283: E971-9.
 7. Kanaya T, Kyo S, Takakura M, Ito H, Namiki M, Inoue M. hTERT is a critical determinant of telomerase activity in renal-cell carcinoma. *Int J Cancer.* 1998 Nov 23;78(5):539-43.
 8. Kaysen JH, Campbell WC, Majewski RR, Goda FO, Navar GL, Lewis FC, Goodwin TJ, Hammond TG. Select de novo gene and protein expression during renal epithelial cell culture in rotating wall vessels is shear stress dependent. *J Membr Biol.* 1999 Mar 1;168(1):77-89.
 9. Keshi H, Sakamoto T, Kawai T, Ohtani K, Katoh T, Jang SJ, Motomura W, Yoshizaki T, Fukuda M, Koyama S, Fukuzawa J, Fukuoh A, Yoshida I, Suzuki Y, Wakamiya N. Identification and characterization of a novel human collectin CL-K1. *Microbiol Immunol.* 2006;50(12):1001-13.
 10. Kiyokawa N, Taguchi T, Mori T, Uchida H, Sato N, Takeda T, Fujimoto J. Induction of apoptosis in normal human renal tubular epithelial cells by Escherichia coli Shiga toxins 1 and 2. *J Infect Dis.* 1998 Jul;178(1):178-84.
 11. Krug AW, Grossmann C, Schuster C, Freudinger R, Mildenberger S, Govindan MV, Gekle M. Aldosterone stimulates epidermal growth factor receptor expression. *J Biol Chem.* 2003 Oct 31;278(44):43060-6.
 12. Kyo S, Takakura M, Taira T, Kanaya T, Itoh H, Yutsudo M, Ariga H, Inoue. Sp1 cooperates with c-Myc to activate transcription of the human telomerase reverse transcriptase gene (hTERT). *Nuc Acids Res.* 2000; 28(3): 669-77.
 13. Loghman-Adham M, Nauli SM, Soto CE, Kariuki B, Zhou J. Immortalized epithelial cells from human autosomal dominant polycystic kidney cysts. *Am J Physiol Renal Physiol.* 2003 Sep;285(3):F397-412.
 14. Loghman-Adham M, Soto CE, Inagami T, Cassis L. The intrarenal renin-angiotensin system in autosomal dominant polycystic kidney disease. *Am J Physiol Renal Physiol.* 2004 Oct;287(4):F775-88.
 15. Matsumura R, Umemiya K, Goto T, Nakazawa T, Kagami M, Tomioka H, Tanabe E, Sugiyama T, Sueishi M. Glandular and extraglandular expression of costimulatory molecules in patients with Sjögren's syndrome. *Ann Rheum Dis.* 2001 May; 60(5):473-82.
 16. Morrissey J, Guo G, Moridaira K, Fitzgerald M, McCracken R, Tolley T, Klahr S. Transforming growth factor-beta induces renal epithelial jagged-1 expression in fibrotic disease. *J Am Soc Nephrol.* 2002 Jun;13(6):1499-508.
 17. Nojima D, Nakajima K, Li LC, Franks J, Ribeiro-Filho L, Ishii N, Dahiya R. CpG methylation of promoter region inactivates E-cadherin gene in renal cell carcinoma. *Mol Carcinog.* 2001 Sep;32(1):19-27.
 18. Prozialeck WC, Edwards JR, Lamar PC. Evaluation of the barrier properties of a commercially available renal epithelial cell system. *Toxicol In Vitro.* 2007 Apr;21(3):533.
 19. Robertson H, Ali S, McDonnell BJ, Burt AD, Kirby JA. Chronic renal allograft dysfunction: the role of T cell-mediated tubular epithelial to mesenchymal cell transition. *J Am Soc Nephrol.* 2004 Feb;15(2):390-7.
 20. Sato F, Abraham JM, Yin J, Kan T, Ito T, Mori Y, Hamilton JP, Jin Z, Cheng Y, Paun B, Berki AT, Wang S, Shimada Y, Meltzer SJ. Polo-like kinase and survivin are esophageal tumor-specific promoters. *Biochem Biophys Res Commun.* 2006 Apr 7;342(2):465-71.
 21. Shen J, Han RG, Karmiol S. Expression of adhesion molecules in cultured human pulmonary microvascular endothelial cells. *Microvasc Res.* 1995; 50: 360-72.
 22. Smith K, Gunaratnam L, Morley M, Franovic A, Mekhail K, Lee S. Silencing of epidermal growth factor receptor suppresses hypoxia-inducible factor-2-driven VHL-/ renal cancer. *Cancer Res.* 2005 Jun 15;65(12):5221-30.
 23. Takakura M, Kyo S, Kanaya T, Hirano H, Takeda J, Yutsudo M, Inoue M. Cloning of

modifications. *Methods Cell Sci.* 2002;24(4):145-53.

- human telomerase catalytic subunit (hTERT) gene promoter and identification of proximal core promoter sequences essential for transcriptional activation in immortalized and cancer cells. *Cancer Res.* 1999 Feb 1;59(3):551-7.
- Takakura M, Kyo S, Sowa Y, Wang Z, Yatabe N, Maida Y, Tanaka M, Inoue M. Telomerase activation by histone deacetylase inhibitor in normal cells. *Nuc Acids Res.* 2001; 29(14): 3006-11.
- Tyler JR, Robertson H, Booth TA, Burt AD, Kirby JA. Chronic allograft nephropathy: intraepithelial signals generated by transforming growth factor-beta and bone morphogenetic protein-7. *Am J Transplant.* 2006 Jun;6(6):1367-76.
- Wilson PD, Hovater JS, Casey CC, Fortenberry JA, Schwiebert EM. ATP release mechanisms in primary cultures of epithelia derived from the cysts of polycystic kidneys. *Am Soc of Nephrol.* 1999; 10.
- Yang Z-Y, Huang Y, Ganesh L, Leung K, Kong W-P, Schqartz O, Subbarao K, Nabel GJ. pH-dependent entry of severe acute respiratory syndrome coronavirus is mediated by the spike glycoprotein and enhanced by dendritic cell transfer through DC-SIGN. *J Virol.* 2004 Jun; 78(11): 5642-50.
- Yatabe N, Kyo S, Kondo S, Kanaya T, Wang Z, Maida Y, Takakura M, Nakamura M, Tanaka M, Inoue M. 2'-5'A antisense therapy directed against human telomerase RNA inhibits telomerase activity and induces apoptosis without telomere impairment in cervical cancer cells. *Cancer Gene Ther.* 2002 Jul;9(7):624-30.
- Hardiek K, Katholi RE, Ramkumar V, Deitrick C. Proximal tubule cell response to radiographic contrast media. *Am J Physiol Renal Physiol.* 2001 Jan;280(1):F61-70.
- Holmén C, Christensson M, Pettersson E, Bratt J, Stjärne P, Karrar A, Sumitran-Holgersson S. Wegener's granulomatosis is associated with organ-specific antiendothelial cell antibodies. *Kidney Int.* 2004 Sep; 66(3):1049-60.
- Keshi H, Sakamoto T, Kawai T, Ohtani K, Katoh T, Jang SJ, Motomura W, Yoshizaki T, Fukuda M, Koyama S, Fukuzawa J, Fukuoh A, Yoshida I, Suzuki Y, Wakamiya N. Identification and characterization of a novel human collectin CL-K1. *Microbiol Immunol.* 2006;50(12):1001-13.
- Lee JE, Park MH, Park JH. The gene expression profile of cyst epithelial cells in autosomal dominant polycystic kidney disease patients. *J Biochem Mol Biol.* 2004 Sep 30;37(5):612-7.
- Maeshima Y, Sudhakar A, Lively JC, Ueki K, Kharbanda S, Kahn CR, Sonenberg N, Hynes RO, Kalluri R. Tumstatin, an endothelial cell-specific inhibitor of protein synthesis. *Sci.* 2002; 295(5552): 140-3.
- Miknyoczki SJ, Jones-Bolin S, Pritchard S, Hunter K, Zhao H, Wan W, Ator M, Bihovsky R, Hudkins R, Chatterjee S, Klein-Szanto A, Dionne C, Ruggeri B. Chemopotentiation of temozolomide, irinotecan, and cisplatin activity by CEP-6800, a poly(ADP-ribose) polymerase inhibitor. *Mol Cancer Ther.* 2003 Apr;2(4):371-82.
- Nukuzuma S, Takasaka T, Zheng HY, Zhong S, Chen Q, Kitamura T, Yogo Y. Subtype I BK polyomavirus strains grow more efficiently in human renal epithelial cells than subtype IV strains. *J Gen Virol.* 2006; 87(Pt 7):1893-901.
- Partha R, Lackey M, Hirsch A, Casscells SW, Conyers JL. Self assembly of amphiphilic C60 fullerene derivatives into nanoscale supramolecular structures. *J Nanobiotechnology.* 2007 Aug 2;5:6.
- Ryan PC, Jakubczak JL, Stewart DA, Hawkins LK, Cheng C, Clarke LM, Ganesh S, Hay C, Huang Y, Kaloss M, Marinov A, Phipps SS, Reddy PS, Shirley PS, Skripchenko Y, Xu L, Yang J, Forry-Schaudies S, Hallenbeck PL. Antitumor efficacy and tumor-selective replication with

HRE

- Aleman MJ, DeYoung MP, Tress M, Keating P, Perry GW, Narayanan R. Inhibition of single minded 2 gene expression mediates tumor-selective apoptosis and differentiation in human colon cancer cells. *PNAS.* 2005 Sep 06; 102(36): 12765-70.
- Chen H, Lo SH. Regulation of tensin-promoted cell migration by its focal adhesion-binding and Src homology 2 domains. *Biochem J.* 2003 Mar 15; 370(Pt 3): 1039-45.
- Elmore E, Siddiqui S, Desai N, Moyer MP, Steele VE, Redpath JL. The human epithelial cell cytotoxicity assay for determining tissue specific toxicity: method

LONZA

- a single intravenous injection of OAS403, an oncolytic adenovirus dependent on two prevalent alterations in human cancer. *Cancer Gene Ther.* 2004 Aug;11(8):555-69.
13. Shin GT, Kim DR, Lim JE, Yim H, Kim H. Upregulation and function of GADD45gamma in unilateral ureteral obstruction. *Kidney Int.* 2008 Jun;73(11):1251-65.
 14. Sugeno Y, Yoshimura A, Yamamura H, Inui K, Morita H, Yamabe H, Ueki N, Ideura T, Takahashi K. Smooth-muscle calponin in mesangial cells: regulation of expression and a role in suppressing glomerulonephritis. *J Am Soc Nephrol.* 2002 Feb; 13(2): 322-31.
 15. Wang Z, Hoy WE. Albuminuria and incident coronary heart disease in Australian Aboriginal people. *Kidney Int.* 2005 Sep; 68(3):1289-93.
 16. Yacoub A, Mitchell C, Brannon J, Rosenberg E, Qiao L, McKinstry R, Linehan WM, Su ZS, Sarkar D, Lebedeva IV, Valerie K, Gopalkrishnan RV, Grant S, Fisher PB, Dent P. MDA-7 (interleukin-24) inhibits the proliferation of renal carcinoma cells and interacts with free radicals to promote cell death and loss of reproductive capacity. *Mol Cancer Ther.* 2003 Jul;2(7):623-32.
 17. Yang Z-Y, Huang Y, Ganesh L, Leung K, Kong W-P, Schqartz O, Subbarao K, Nabel GJ. pH-dependent entry of sever acute respiratory syndrome coronavirus is mediated by the spike glycoprotein and enhanced by dendritic cell transfer through DC-SIGN. *J Virol.* 2004 Jun; 78(11): 5642-50.
- NHMC⁺**
1. Amemiya T, Sasamura H, Mifune M, Kitamura Y, Hirahashi J, Hayashi M, Saruta T. Vascular endothelial growth factor activates MAP kinase and enhances collagen synthesis in human mesangial cells. *Kidney Int.* 1999 Dec;56(6):2055-63.
 2. Banba N, Nakamura T, Matsumura M, Kuroda H, Hattori Y, Kasai K. Possible relationship of monocyte chemoattractant protein-1 with diabetic nephropathy. *Kidney Int.* 2000 Aug;58(2):684-90.
 3. Basler CF, García-Sastre A, Palese P. A novel paramyxovirus? *Emerg Infect Dis.* 2005 Jan;11(1):108-12.
 4. Bassa BV, Noh JW, Ganji SH, Shin MK, Roh DD, Kamanna VS. Lysophosphatidylcholine stimulates EGF receptor activation and mesangial cell proliferation: regulatory role of Src and PKC. *Biochim Biophys Acta.* 2007 Nov;1771(11):1364-71.
 5. Beqaj S, Shah AM, Ryan JM. Identification of cells responsible for urokinase-type plasminogen activator synthesis and secretion in human diploid kidney cell cultures. *In Vitro Cell Dev Biol Anim.* 2004 Mar-Apr;40(3-4):102-7.
 6. Black SA Jr, Palamakumbura AH, Stan M, Trackman PC. Tissue-specific mechanisms for CCN2/CTGF persistence in fibrotic gingiva: interactions between cAMP and MAPK signaling pathways, and prostaglandin E2-EP3 receptor mediated activation of the c-JUN N-terminal kinase. *J Biol Chem.* 2007 May 25;282(21):15416-29.
 7. Bobst SM, Day MC, Gilstrap LC 3rd, Xia Y, Kellems RE. Maternal autoantibodies from preeclamptic patients activate angiotensin receptors on human mesangial cells and induce interleukin-6 and plasminogen activator inhibitor-1 secretion. *Am J Hypertens.* 2005 Mar;18(3):330-6.
 8. Bolick DT, Hatley ME, Srinivasan S, Hedrick CC, Nadler JL. Lisofylline, a novel antiinflammatory compound, protects mesangial cells from hyperglycemia- and angiotensin II-mediated extracellular matrix deposition. *Endocrinology.* 2003 Dec;144(12):5227-31.
 9. Chaudhary A, Mishra A, Sethi S. Oxidized omega-3 fatty acids inhibit pro-inflammatory responses in glomerular endothelial cells. *Nephron Exp Nephrol.* 2004;97(4):e136-45.
 10. Chromek M, Tullus K, Hertting O, Jaremk G, Khalil A, Li Y-H, Brauner A. Matrix metalloproteinase-9 and tissue inhibitor of metalloproteinases-1 in acute pyelonephritis and renal scarring. *Pediat Res.* 2003; 53(4): 698-705.
 11. Coletta I, Soldo L, Polentarutti N, Mancini F, Guglielmotti A, Pinza M, Mantovani A, Milanese C. Selective induction of MCP-1 in human mesangial cells by the IL-6/SIL-6R complex. *Exp Nephrol.* 2000; 8(1): 37-43.
 12. Du J, Ding M, Sours-Brothers S, Graham S, Ma R. Mediation of angiotensin II-induced Ca²⁺ signaling by polycystin 2 in glomerular mesangial cells. *Am J Physiol Renal Physiol.* 2008 Apr;294(4):F909-18.+
 13. Dubey RK, Gillespie DG, Keller PJ, Imthurn B, Zacharia LC, Jackson EK. Role of methoxyestradiols in the growth inhibitory

- effects of estradiol on human glomerular mesangial cells. *Hypertension*. 2002 Feb;39(2 Pt 2):418-24.
14. Endo S, Hiramatsu N, Hayakawa K, Okamura M, Kasai A, Tagawa Y, Sawada N, Yao J, Kitamura M. Geranylgeranylacetone, an inducer of the 70-kDa heat shock protein (HSP70), elicits unfolded protein response and coordinates cellular fate independently of HSP70. *Mol Pharmacol*. 2007 Nov;72(5):1337-48.
 15. Eto S, Isome M, Sano H, Fukuda Y, Kawasaki Y, Suzuki J, Igarashi K, Satriano J, Suzuki H. Agmatine suppresses mesangial cell proliferation by modulating polyamine metabolism. *Tohoku J Exp Med*. 2006 Oct;210(2):145-51.
 16. Etoh T, Inoguchi T, Kakimoto M, Sonoda N, Kobayashi K, Kuroda J, Sumimoto H, Nawata H. Increased expression of NAD(P)H oxidase subunits, NOX4 and p22phox, in the kidney of streptozotocin-induced diabetic rats and its reversibility by interventive insulin treatment. *Diabetologia*. 2003 Oct;46(10):1428-37.
 17. Fan YP, Weiss RH. Exogenous attenuation of p21(Waf1/Cip1) decreases mesangial cell hypertrophy as a result of hyperglycemia and IGF-1. *J Am Soc Nephrol*. 2004 Mar;15(3):575-84.
 18. Furlong F, Crean J, Thornton L, O'Leary R, Murphy M, Martin F. Dysregulated intracellular signaling impairs CTGF-stimulated responses in human mesangial cells exposed to high extracellular glucose. *Am J Physiol Renal Physiol*. 2007 Jun;292(6):F1691-700.
 19. Harper L, Kashiwagi Y, Pusey CD, Hendry BM, Domin J. Platelet-derived growth factor reorganizes the actin cytoskeleton through 3-phosphoinositide-dependent and 3-phosphoinositide-independent mechanisms in human mesangial cells. *Nephron Physiol*. 2007;107(2):p45-56.
 20. Huang Y, Wongamorntham S, Kasting J, McQuillan D, Owens RT, Yu L, Noble NA, Border W. Renin increases mesangial cell transforming growth factor-beta1 and matrix proteins through receptor-mediated, angiotensin II-independent mechanisms. *Kidney Int*. 2006 Jan;69(1):105-13.
 21. Hughes AK, Stricklett PK, Schmid D, Kohan DE. Cytotoxic effect of Shiga toxin-1 on human glomerular epithelial cells. *Kidney Int*. 2000 Jun;57(6):2350-9.
 22. Jung EM, Park JW, Choi KS, Park JW, Lee HI, Lee KS, Kwon TK. Curcumin sensitizes tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-mediated apoptosis through CHOP-independent DR5 upregulation. *Carcinogenesis*. 2006 Oct;27(10):2008-17.
 23. Kanasaki K, Koya D, Sugimoto T, Isono M, Kashiwagi A, Haneda M. N-acetyl-seryl-aspartyl-lysyl-proline inhibits TGF- β -mediated plasminogen activator inhibitor-1 expression via inhibition of Smad pathway in human Mesangial cells. *J Am Soc Nephrol*. 2003; 14: 863-72.
 24. Kapasi AA, Fan S, Singhal PC. Role of 14-3-3epsilon, c-Myc/Max, and Akt phosphorylation in HIV-1 gp 120-induced mesangial cell proliferation. *Am J Physiol Renal Physiol*. 2001 Feb;280(2):F333-42.
 25. Kapasi AA, Fan S, Singhal PC. Scatter factor mitigates HIV-1 gp120-induced human mesangial cell injury. *Nephron Exp Nephrol*. 2006;103(3):e103-8.
 26. Keshi H, Sakamoto T, Kawai T, Ohtani K, Katoh T, Jang SJ, Motomura W, Yoshizaki T, Fukuda M, Koyama S, Fukuzawa J, Fukuoh A, Yoshida I, Suzuki Y, Wakamiya N. Identification and characterization of a novel human collectin CL-K1. *Microbiol Immunol*. 2006;50(12):1001-13.
 27. Khwaja A, Dockrell ME, Hendry BM, Sharpe CC. Prenylation is not necessary for endogenous Ras activation in non-malignant cells. *J Cell Biochem*. 2006 Feb 1;97(2):412-22.
 28. Kitahara T, Hiromura K, Ikeuchi H, Yamashita S, Kobayashi S, Kuroiwa T, Kaneko Y, Ueki K, Nojima Y. Mesangial cells stimulate differentiation of endothelial cells to form capillary-like networks in a three-dimensional culture system. *Nephrol Dial Transplant*. 2005 Jan;20(1):42-9.
 29. Kobayashi T, Furukawa Y, Kikuchi J, Ito C, Miyata Y, Muto S, Tanaka A, Kusano E. Transactivation of RON receptor tyrosine kinase by interaction with PDGF receptor beta during steady-state growth of human mesangial cells. *Kidney Int*. 2009 Jun;75(11):1173-83.
 30. Koyama T, Hora K, Ako S, Kamijo Y, Higuchi M. Expression of hepatocyte growth factor is induced by the interaction between human mesangial cells and monocytes. *Nephron Exp Nephrol*. 2003; 94(4): e146-53.

31. Lee TJ, Jung EM, Lee JT, Kim S, Park JW, Choi KS, Kwon TK. Mithramycin A sensitizes cancer cells to TRAIL-mediated apoptosis by down-regulation of XIAP gene promoter through Sp1 sites. *Mol Cancer Ther.* 2006 Nov; 5(11):2737-46.
32. Lee TJ, Um HJ, Min do S, Park JW, Choi KS, Kwon TK. Withaferin A sensitizes TRAIL-induced apoptosis through reactive oxygen species-mediated up-regulation of death receptor 5 and down-regulation of c-FLIP. *Free Radic Biol Med.* 2009 Jun 15;46(12):1639-49.
33. Liang XB, Zhang H, Zhou AY, Wang HY. Screening for functional genes related to a novel gene, AngRem104, in human mesangial cells by cDNA microarray. *Biotechnol Lett.* 2003 Jan;25(2):139-42.
34. Lim JH, Park JW, Choi KS, Park YB, Kwon TK. Rottlerin induces apoptosis via death receptor 5 (DR5) upregulation through CHOP-dependent and PKC delta-independent mechanism in human malignant tumor cells. *Carcinogenesis.* 2009 May;30(5):729-36.
35. Lipkowitz MS, Hanss B, Tulchin N, Wilson PD, Langer JC, Ross MD, Kurtzman GJ, Klotman PE, Klotman ME. Transduction of renal cell *sin vitro* and *in vivo* by adeno-associated virus gene therapy vectors. 1999; 10: 1908-15.
36. Luo P, Tan Z, Zhang Z, Li H, Mo Z. Inhibitory effects of salvianolic acid B on the high glucose-induced mesangial proliferation via NF-kappaB-dependent pathway. *Biol Pharm Bull.* 2008 Jul;31(7):1381-6.
37. Matsui T, Yamagishi S, Takeuchi M, Ueda S, Fukami K, Okuda S. Nifedipine, a calcium channel blocker, inhibits advanced glycation end product (AGE)-elicited mesangial cell damage by suppressing AGE receptor (RAGE) expression via peroxisome proliferator-activated receptor-gamma activation. *Biochem Biophys Res Commun.* 2009 Jul 24;385(2):269-72.
38. Mattana J, Kochlatyi S, Gibbons N. Metal-catalyzed oxidation of extracellular matrix proteins promotes human mesangial cell apoptosis and is associated with enhanced expression of Bax and caspase activation. *Biochem Biophys Res Commun.* 2002 Apr 5;292(3):652-8.
39. Melnyk RA, Tam J, Boie Y, Kennedy BP, Percival MD. Renin and prorenin activate pathways implicated in organ damage in human mesangial cells independent of angiotensin II production. *Am J Nephrol.* 2009;30(3):232-43.
40. Mishra R, Emancipator SN, Kern T, Simonson MS. High glucose evokes an intrinsic proapoptotic signaling pathway in mesangial cells. *Kidney Int.* 2005 Jan;67(1):82-93.
41. Mishra R, Simonson MS. Oleate induces a myofibroblast-like phenotype in mesangial cells. *Arterioscler Thromb Vasc Biol.* 2008 Mar;28(3):541-7.
42. Nakamura H, Isaka Y, Tsujie M, Akagi Y, Sudo T, Ohno N, Imai E, Hori M. Electroporation-mediated PDGF receptor-IgG chimera gene transfer ameliorates experimental glomerulonephritis. *Kidney Int.* 2001 Jun;59(6):2134-45.
43. Negishi M, Shimizu H, Okada S, Kuwabara A, Okajima F, Mori M. 9HODE stimulates cell proliferation and extracellular matrix synthesis in human mesangial cells via PPARgamma. *Exp Biol Med (Maywood).* 2004 Nov;229(10):1053-60.
44. Nishikawa T, Suematsu S, Saito J, Soyama A, Ito H, Kino T, Chrousos G. Human renal mesangial cells produce aldosterone in response to low-density lipoprotein (LDL). *J Steroid Biochem Mol Biol.* 2005 Aug;96(3-4):309-16.
45. Oortwijn BD, van der Boog PJ, Roos A, van der Geest RN, de Fijter JW, Daha MR, van Kooten C. A pathogenic role for secretory IgA in IgA nephropathy. *Kidney Int.* 2006 Apr; 69(7):1131-8.
46. Prabhakar SS. Inhibition of Mesangial iNOS by reduced extracellular pH is associated with uncoupling of NADPH oxidation. *Kidney Int.* 2002; 61: 2015-24.
47. Russell WJ, Cardelli J, Harris E, Baier RJ, Herrera GA. Monoclonal light chain--mesangial cell interactions: early signaling events and subsequent pathologic effects. *Lab Invest.* 2001 May;81(5):689-703.
48. Sawai K, Mukoyama M, Mori K, Kasahara M, Koshikawa M, Yokoi H, Yoshioka T, Ogawa Y, Sugawara A, Nishiyama H, Yamada S, Kuwahara T, Saleem MA, Shiota K, Ogawa O, Miyazato M, Kangawa K, Nakao K. Expression of CCN1 (CYR61) in developing, normal, and diseased human kidney. *Am J Physiol Renal Physiol.* 2007 Oct;293(4):F1363-72.

49. Schwerdt G, Holzinger H, Königs M, Humpf HU, Gekle M. Effect of ochratoxin A on cell survival and collagen homeostasis in human mesangial cells in primary culture. *Food Chem Toxicol.* 2009 Jan;47(1):209-13.
50. Shushakova N, Tkachuk N, Dangers M, Tkachuk S, Park JK, Hashimoto K, Haller H, Dumler I. Urokinase-induced activation of the gp130/Tyk2/Stat3 pathway mediates a pro-inflammatory effect in human mesangial cells via expression of the anaphylatoxin C5a receptor. *J Cell Sci.* 2005 Jun 15;118(Pt 12):2743-53.
51. Singh R, Browning JL, Abi-Habib R, Wong K, Williams SA, Merchant R, Denmeade SR, Buckley TJ, Frankel AE. Recombinant prostate-specific antigen proaerolysin shows selective protease sensitivity and cell cytotoxicity. *Anticancer Drugs.* 2007 Aug;18(7):809-16.
52. Sugenoya Y, Yoshimura A, Yamaura H, Inui K, Morita H, Yamabe H, Ueki N, Ideura T, Takahashi K. Smooth-muscle calponin in mesangial cells: regulation of expression and a role in suppressing glomerulonephritis. *J Am Soc Nephrol.* 2002; 13: 322-31.
53. Sweeney EA, Igarashi Y. Increases in ceramide levels in normal human mesangial cells subjected to different cellular stresses result from changes in distinct enzyme activities and can influence cellular responses to other stimuli. *J Biochem.* 1999 Apr;125(4):737-45.
54. Tahara A, Tsukada J, Tomura Y, Momose K, Suzuki T, Yatsu T, Shibasaki M. Effects of YM218, a nonpeptide vasopressin V(1A) receptor-selective antagonist, on vasopressin-induced growth responses in human mesangial cells. *Eur J Pharmacol.* 2006 May 24;538(1-3):32-8.
55. Xing Y, Ganji SH, Noh JW, Kamanna VS. Cell density-dependent expression of EDG family receptors and mesangial cell proliferation: role in lysophosphatidic acid-mediated cell growth. *Am J Physiol Renal Physiol.* 2004 Dec;287(6):F1250-7.
56. Yasuoka H, Yoshizumi M, Inui D, Okishima N, Houchi H, Kirima K, Oshita S, Kido H, Tamaki T. Effect of endothelin-1(1-31) on intracellular free calcium in cultured human mesangial cells. *Life Sci.* 1999;65(22):PL267-72.
57. Yu HY, Inoguchi T, Nakayama M, Tsubouchi H, Sato N, Sonoda N, Sasaki S, Kobayashi K, Nawata H. Statin attenuates high glucose-induced and angiotensin II-induced MAP kinase activity through inhibition of NAD(P)H oxidase activity in cultured mesangial cells. *Med Chem.* 2005 Sep;1(5):461-6.
58. Zernecke A, Weber KS, Erwig LP, Kluth DC, Schröppel B, Rees AJ, Weber C. Combinatorial model of chemokine involvement in glomerular monocyte recruitment: role of CXC chemokine receptor 2 in infiltration during nephrotoxic nephritis. *J Immunol.* 2001 May 1;166(9):5755-62.
59. Zhang Y, Chen K, Guo L, Wu C. Characterization of PINCH-2, a new focal adhesion protein that regulates the PINCH-1-ILK interaction, cell spreading, and migration. *J Biol Chem.* 2002 Oct 11;277(41):38328-38.
60. Zhang Y, Yasumoto Y, Ikeda T, Takenouchi S, Sogabe A, Nosaki T, Che X, Zheng C, Haraguchi M, Akiyama S, Tsubouchi H. Apolipoprotein E regulates primary cultured human mesangial cell proliferation. *Nephron Exp Nephrol.* 2006;102(2):e62-70.
61. Zhang Y, Yasumoto Y, Mei C, Arima T. Vitamin E inhibits proliferation of primary cultured human mesangial and endothelial cells. *Nephron.* 2001 Nov; 89(3):291-6.
62. Zhao Z, Weinstein E, Tuzova M, Davidson A, Mundel P, Marambio P, Puttermann C. Cross-reactivity of human lupus anti-DNA antibodies with alpha-actinin and nephritogenic potential. *Arthritis Rheum.* 2005 Feb;52(2):522-30.

Media

1. Kunduzova OR, Escourrou G, De La Farge F, Salvayre R, Séguélas M-H, Leducq N, Bono F, Herbert J-M, Parini A. Involvement of peripheral benzodiazepine receptor in the oxidative stress, death-signaling pathways, and renal injury induced by ischemia-reperfusion. *J Am Soc Nephrol.* 2004; 15: 2152-60.
2. Liu XH, Aigner A, Wellstein A, Ray PE. Up-regulation of a fibroblast growth factor binding protein in children with renal diseases. *Kidney Int.* 2001 May;59(5):1717-28.
3. Phillips JL, Ghadimi BM, Wangsa D, Padilla-Nash H, Worrell R, Hewitt S, Walther M, Linehan WM, Klausner RD, Ried T. Molecular cytogenetic characterization of early and late renal cell carcinomas in von

Lonza

Hippel-Lindau disease. Genes
Chromosomes Cancer. 2001 May;31(1):1-9.

* References not specifically citing the use of Lonza cells, media,
or reagents in their research.

+ Denotes sections containing only the articles published within
the last ten years.