

## Product References

### Clonetics® Skeletal Muscle Cells

---

#### Cells

#### HSMC

1. Ates K, Yang SY, Orrell RW, Sinanan AC, Simons P, Solomon A, Beech S, Goldspink G, Lewis MP. The IGF-I splice variant MGF increases progenitor cells in ALS, dystrophic, and normal muscle. *FEBS Lett.* 2007 Jun 12;581(14):2727-32.
2. Baffour R, Pakala R, Hellinga D, Joner M, Okubagzi P, Epstein SE, Waksman R. Bone marrow-derived stem cell interactions with adult cardiomyocytes and skeletal myoblasts in vitro. *Cardiovasc Revasc Med.* 2006 Oct-Dec;7(4):222-30.
3. Buas MF, Kabak S, Kadesch T: Inhibition of myogenesis by Notch: evidence for multiple pathways. *J Cell Physiol.* 2009 Jan;218(1):84-93.
4. Chowdhury SR, Muneyuki Y, Takezawa Y, Kino-Oka M, Saito A, Sawa Y, Taya M. Synergic stimulation of laminin and epidermal growth factor facilitates the myoblast growth through promoting migration. *J Biosci Bioeng.* 2009 Aug;108(2):174-7.
5. Frost RA, Lang CH, Gelato MC. Transient exposure of human myoblasts to tumor necrosis factor- $\alpha$  inhibits serum and insulin-like growth factor-I stimulated protein synthesis. *Endocrinol.* 1997; 138: 4153-9.
6. Girgenrath M, Weng S, Kostek CA, Browning B, Wang M, Brown SA, Winkles JA, Michaelson JS, Allaire N, Schneider P, Scott ML, Hsu YM, Yagita H, Flavell RA, Miller JB, Burkly LC, Zheng TS. TWEAK, via its receptor Fn14, is a novel regulator of mesenchymal progenitor cells and skeletal muscle regeneration. *EMBO J.* 2006; 25(24): 5826-39.
7. Grozovsky R, Ribich S, Rosene ML, Mulcahey MA, Huang SA, Patti ME, Bianco AC, Kim BW. Type 2 deiodinase expression is induced by peroxisomal proliferator-activated receptor- $\gamma$  agonists in skeletal myocytes. *Endocrinology.* 2009 Apr;150(4):1976-83.
8. Huang SA, Mulcahey MA, Crescenzi A, Chung M, Kim BW, Barnes C, Kuijt W, Turano H, Harney J, Larsen PR. Transforming growth factor- $\beta$  promotes inactivation of extracellular thyroid hormones via transcriptional stimulation of type 3 iodothyronine deiodinase. *Mol Endocrinol.* 2005 Dec;19(12):3126-36.
9. Kim DJ, Oh B, Kim YY. Splicing factor ASF/SF2 and transcription factor PPAR- $\gamma$  cooperate to directly regulate transcription of uncoupling protein-3. *Biochem Biophys Res Commun.* 2009 Jan 23;378(4):877-82.
10. Kino-Oka M, Chowdhury SR, Muneyuki Y, Manabe M, Saito A, Sawa Y, Taya M. Automating the expansion process of human skeletal muscle myoblasts with suppression of myotube formation. *Tissue Eng Part C Methods.* 2009 Mar 13.
11. Lee JH, Bugarija B, Millan EJ, Walton NM, Gaetz J, Fernandes CJ, Yu WH, Mekel-Bobrov N, Vallender TW, Snyder GE, Xiang AP, Lahn BT. Systematic identification of cis-silenced genes by trans complementation. *Hum Mol Genet.* 2009 Mar 1;18(5):835-46.
12. Lee JH, Gaetz J, Bugarija B, Fernandes CJ, Snyder GE, Bush EC, Lahn BT. Chromatin analysis of occluded genes. *Hum Mol Genet.* 2009 Jul 15;18(14):2567-74.
13. Linardic CM, Downie DL, Qualman S, Bentley RC, Counter CM. Genetic modeling of human rhabdomyosarcoma. *Canc Res.* 2005; 65: 4490-5.
14. Liu S-H, Towler MC, Chen E, Chen C-Y, Song W, Apodaca G, Brodsky FM. A novel

- clathrin homolog that co-distributes with cytoskeletal components functions in the trans-Golgi network. *EMBO J.* 2001; 20 (1 & 2): 272-84.
15. Marquette ML, Byerly D, Sognier M. A novel in vitro three-dimensional skeletal muscle model. *In Vitro Cell Dev Biol Anim.* 2007 Jul-Aug;43(7):255-63.
  16. McGuinness R. Label-free cell-based assays bring greater biorelevance to early drug discovery. *Assay Drug Dev Technol.* 2009 Apr;7(2):191-3.
  17. Menezes KM, Mok HS, Barry MA. Increased transduction of skeletal muscle cells by fibroblast growth factor-modified adenoviral vectors. *Human Gene Ther.* 2006 Mar; 17(3): 314-20.
  18. Naini S, Etheridge KT, Adam SJ, Qualman SJ, Bentley RC, Counter CM, Linardic CM. Defining the cooperative genetic changes that temporally drive alveolar rhabdomyosarcoma. *Cancer Res.* 2008 Dec 1;68(23):9583-8.
  19. Nouette-Gaulain K, Bellance N, Prévost B, Passerieux E, Pertuiset C, Galbes O, Smolkova K, Masson F, Miraux S, Delage JP, Letellier T, Rossignol R, Capdevila X, Sztark F: Erythropoietin protects against local anesthetic myotoxicity during continuous regional analgesia. *Anesthesiology.* 2009 Mar;110(3):648-59.
  20. Padmalayam I, Hasham S, Saxena U, Pillarisetti S. Lipoic acid synthase (LASy): a novel role in inflammation, mitochondrial function, and insulin resistance. *Diabetes.* 2009 Mar;58(3):600-8.
  21. Qi JS, Minor LK, Smith C, Hu B, Yang J, Andrade-Gordon P, Damiano B. Characterization of functional urotensin II receptors in human skeletal muscle myoblasts: comparison with angiotensin II receptors. *Peptides.* 2005 Apr;26(4):683-90.
  22. Saitoh A, Haas RH, Naviaux RK, Salva NG, Wong JK, Spector SA. Impact of nucleoside reverse transcriptase inhibitors on mitochondrial DNA and RNA in human skeletal muscle cells. *Antimicrob Agents Chemother.* 2008 Aug;52(8):2825-30.
  23. Senetar MA, Moncman CL, McCann RO. Talin2 is induced during striated muscle differentiation and is targeted to stable adhesion complexes in mature muscle. *Cell Motil Cytoskeleton.* 2007 Mar;64(3):157-73.
  24. Smith KP, Byron M, Johnson C, Xing Y, Lawrence JB. Defining early steps in mRNA transport: mutant mRNA in myotonic dystrophy type I is blocked at entry into SC-35 domains. *J Cell Biol.* 2007 Sep 10;178(6):951-64.
  25. Su Y, Vanderlaag K, Ireland C, Ortiz J, Grage H, Safe S, Frankel AE. 1,1-Bis(3'-indolyl)-1-(p-biphenyl)methane inhibits basal-like breast cancer growth in athymic nude mice. *Breast Cancer Res.* 2007;9(4):R56.
  26. Watanabe M, Houten SM, Matakai C, Christoffolete MA, Kim BW, Sato H, Messaddeq N, Harney JW, Ezaki O, Kodama T, Schoonjans K, Bianco AC, Auwerx J. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature.* 2006 Jan 26;439(7075):484-9.
  27. Zhao W, Pan J, Wang X, Wu Y, Bauman WA, Cardozo CP. Expression of the muscle atrophy factor muscle atrophy F-box is suppressed by testosterone. *Endocrinology.* 2008 Nov;149(11):5449-60.

## SkMC

1. Anderson K, Potter A, Baban D, Davies KE. Protein expression changes in spinal muscular atrophy revealed with a novel antibody array technology. *Brain.* 2003 Jul7; 126: 2052-64.
2. Benayoun B, Baghdiguian S, Lajmanovich A, Bartoli M, Daniele N, Gicquel E, Bourg N, Raynaud F, Pasquier MA, Suel L, Lochmuller H, Lefranc G, Richard I. NF-kappaB-dependent expression of the antiapoptotic factor c-FLIP is regulated by calpain 3, the protein involved in limb-girdle muscular dystrophy type 2A. *FASEB J.* 2008 May;22(5):1521-9.
3. Binley K, Iqbal S, Kingsman A, Kingsman S, Naylor S. An adenoviral vector regulated by hypoxia for the treatment of ischaemic disease and cancer. *Gene Ther.* 1999; 6(10):1721-7.
4. Birkus G, Gibbs CS, Cihlar T. Comparative effects of adefovir and selected nucleosides inhibitors of hepatitis B virus DNA polymerase on mitochondrial DNA in liver and skeletal muscle cells. *J Viral Hepat.* 2003 Jan; 10 (1): 50-4.
5. Bryant AE, Bayer CR, Huntington JD, Stevens DL. Group A streptococcal myonecrosis: increased vimentin expression after skeletal-muscle injury mediates the

- binding of *Streptococcus pyogenes*. *J Infect Dis*. 2006 Jun 15;193(12):1685-92.
6. Bunas MF, Kabak S, Kadesch T: Inhibition of myogenesis by Notch: evidence for multiple pathways. *J Cell Physiol*. 2009 Jan;218(1):84-93.
  7. Dijkmans J, Xu J, Masure S, Dhanaraj S, Gosiewska A, Geesin J, Sprengel J, Harris S, Verhasselt P, Gordon R, Yon J. Characterization of platelet-derived growth factor-C (PDGF-C): expression in normal and tumor cells, biological activity and chromosomal localization. *Int J Biochem Cell Biol*. 2002 Apr;34(4):414-26.
  8. Fougousse F, Durand M, Lopez S, Suel L, Demignon J, Thornton C, Ozaki H, Kawakami K, Barbet JS, Maire P. Six and eya expression during human somitogenesis and MyoD gene family activation. *J Musc Res Cell Motil*. 2002 Apr; 23(3): 255-64.
  9. Fulco M, Schiltz RL, Iezzi S, King MT, Zhao P, Kashiwaya Y, Hoffman E, Veech RL, Sartorelli V. Sir2 regulates skeletal muscle differentiation as a potential sensor of the redox state. *Mol Cell*. 2003 Jul;12(1):51-62.
  10. Gramolini AO, Angus LM, Schaeffer L, Burton EA, Tinsley JM, Davies KE, Changeux JP, Jasmin BJ. Induction of utrophin gene expression by heregulin in skeletal muscle cells: role of the N-box motif and GA binding protein. *Proc Natl Acad Sci U S A*. 1999 Mar 16;96(6):3223-7.
  11. Grozovsky R, Ribich S, Rosene ML, Mulcahey MA, Huang SA, Patti ME, Bianco AC, Kim BW: Type 2 deiodinase expression is induced by peroxisomal proliferator-activated receptor-gamma agonists in skeletal myocytes. *Endocrinology*. 2009 Apr;150(4):1976-83.
  12. Hong-Brown LQ, Frost RA, Lang CH. Alcohol impairs protein synthesis and degradation in cultured skeletal muscle cells. *Alcohol Clin Exp Res*. 2001 Sep;25(9):1373-82.
  13. Hu J, Higuchi I, Yoshida Y, Shiraishi T, Osame M. Expression of midkine in regenerating skeletal muscle fibers and cultured myoblasts of human skeletal muscle. *Eur Neurol*. 2002; 47: 20-5.
  14. Krishnamurthy S, Takimoto T, Scroggs RA, Portner A. Differentially regulated interferon response determines the outcome of Newcastle disease virus infection in normal and tumor cell lines. *J Virol*. 2006; 80(11):5145-55.
  15. Linardic CM, Naini S, Herndon JE 2nd, Kesslerwan C, Qualman SJ, Counter CM. The PAX3-FKHR fusion gene of rhabdomyosarcoma cooperates with loss of p16INK4A to promote bypass of cellular senescence. *Cancer Res*. 2007 Jul 15;67(14):6691-9.
  16. Liu SH, Towler MC, Chen E, Chen CY, Song W, Apodaca G, Brodsky FM. A novel clathrin homolog that co-distributes with cytoskeletal components functions in the trans-Golgi network. *EMBO J*. 2001 Jan 15;20(1-2):272-84.
  17. Lo WR, Rowlette LL, Caballero M, Yang P, Hernandez MR, Borrás T. Tissue differential microarray analysis of dexamthasone induction reveals potential mechanisms of steroid glaucoma. *Investig Ophthal & Visual Sci*. 2003; 44: 473-85.
  18. Minnich A, Tian N, Byan L, Bilder G. A potent PPARalpha agonist stimulates mitochondrial fatty acid beta-oxidation in liver and skeletal muscle. *Am J Physiol Endocrinol Metab*. 2001 Feb;280(2):E270-9.
  19. Mizuno H, Zuk PA, Zhu M, Lorenz HP, Benhaim P, Hedrick MH. Myogenic differentiation by human processed lipoaspirate cells. *Plast Reconstr Surg*. 2002 Jan;109(1):199-209.
  20. Molina RP, Matukonis M, Paszkiet B, Zhang J, Kaleko M, Luo T. Mapping of the bovine immunodeficiency virus packaging signal and RRE and incorporation into a minimal gene transfer vector. *Virol*. 2002; 301: 10-23.
  21. Montano M, Flanagan JN, Jiang L, Sebastiani P, Rarick M, LeBrasseur NK, Morris CA, Jasuja R, Bhasin S. Transcriptional profiling of testosterone-regulated genes in the skeletal muscle of human immunodeficiency virus-infected men experiencing weight loss. *J Clin Endocrinol Metab*. 2007 Jul;92(7):2793-802.
  22. Nagaraju K, Casciola-Rosen L, Rosen A, Thompson C, Loeffler L, Parker T, Danning C, Rochon PJ, Gillespie J, Plotz P. The inhibition of apoptosis in myositis and in normal muscle cells. *J Immunol*. 2000 May 15;164(10):5459-65.
  23. Nagaraju K, Raben N, Merritt G, Loeffler L, Kirk K, Plotz P. A variety of cytokines and immunologically relevant surface molecules are expressed by normal human skeletal muscle cells under proinflammatory stimuli. *Clin Exp Immunol*. 1998 Sep;113(3):407-14.
  24. Newton K, French DM, Yan M, Frantz GD, Dixit VM. Myodegeneration in EDA-A2

- transgenic mice is prevented by XEDAR deficiency. *Mol Cell Biol.* 2004 Feb;24(4):1608-13.
25. Philips AV, Timchenko LT, Cooper TA. Disruption of splicing regulated by a CUG-binding protein in myotonic dystrophy. *Science.* 1998 May 1;280(5364):737-41.
  26. Saravia-Otten P, Frisan T, Thelestam M, Gutiérrez JM. Membrane independent activation of fibroblast proMMP-2 by snake venom: novel roles for venom proteinases. *Toxicon.* 2004 Dec 1;44(7):749-64.
  27. Shearer BG, Steger DJ, Way JM, Stanley TB, Lobe DC, Grillot DA, Iannone MA, Lazar MA, Willson TM, Billin AN. Identification and characterization of a selective peroxisome proliferator-activated receptor beta/delta (NR1C2) antagonist. *Mol Endocrinol.* 2008 Feb;22(2):523-9.
  28. Sinha-Hikim I, Taylor WE, Gonzalez-Cadavid NF, Zheng W, Bhasin S. Androgen receptor in human skeletal muscle and cultured muscle satellite cells: up-regulation by androgen treatment. *J Clin Endocrinol & Metabol.* 2004; 89(10): 5245-55.
  29. Solanes G, Pedraza N, Iglesias R, Giralt M, Villarrojo F. Functional relationship between MyoD and peroxisome proliferator-activated receptor-dependent regulatory pathways in the control of the human uncoupling protein-3 gene transcription. *Mol Endocrin.* 2003; 17(10): 1944-58.
  30. Taniyama Y, Tachibana K, Hiraoka K, Aoki M, Yamamoto S, Matsumoto K, Nakamura T, Ogihara T, Kaneda Y, Morishita R. Development of safe and efficient novel nonviral gene transfer using ultrasound: enhancement of transfection efficiency of naked plasmid DNA in skeletal muscle. *Gene Ther.* 2002 Mar;9(6):372-80.
  31. Tucker DE, Stewart A, Nallan L, Bendale P, Ghomashchi F, Gelb MH, Leslie CC. Group IVC cytosolic phospholipase A<sub>2γ</sub> is farnesylated and palmitoylated in mammalian cells. *J Lipid Res.* 2005; 46: 2122-33.
  32. Walczak H, Miller RE, Ariail K, Gliniak B, Griffith TS, Kubin M, Chin W, Jones J, Woodward A, Le T, Smith C, Smolak P, Goodwin RG, Rauch CT, Schuh JC, Lynch DH. Tumoricidal activity of tumor necrosis factor-related apoptosis-inducing ligand in vivo. *Nat Med.* 1999; 5(2):157-63.
  33. Young HE, Steele TA, Bray RA, Detmer K, Blake LW, Lucas PW, Black AC Jr. Human pluripotent and progenitor cells display cell surface cluster differentiation markers CD10, CD13, CD56, and MHC class-I. *Proc Soc Exp Biol Med.* 1999 May;221(1):63-71.
  34. Young HE, Steele TA, Bray RA, Hudson J, Floyd JA, Hawkins K, Thomas K, Austin T, Edwards C, Cuzzourt J, Duenzl M, Lucas PA, Black AC Jr. Human reserve pluripotent mesenchymal stem cells are present in the connective tissues of skeletal muscle and dermis derived from fetal, adult, and geriatric donors. *Anat Rec.* 2001 Sep 1;264(1):51-62.

## Media

1. Bertacca A, Ciccarone A, Cecchetti P, Vianello B, Laurenza I, Maffei M, Chiellini C, Del Prato S, Benzi L. Continually high insulin levels impair Akt phosphorylation and glucose transport in human myoblasts. *Metabolism.* 2005; 54(12):1687-93.
2. Kuçi S, Kuçi Z, Schmid S, Seitz G, Müller I, Dufke A, Leimig T, Murti G, Jurecic R, Schumm M, Lang P, Bruchelt G, Bader P, Klingebiel T, Niethammer D, Handgretinger R. Efficient in vitro generation of adult multipotent cells from mobilized peripheral blood CD133+ cells. *Cell Prolif.* 2008 Feb;41(1):12-27.
3. McCann JA, Muro EM, Palmer C, Palidwor G, Porter CJ, Andrade-Navarro MA, Rudnicki MA. ChIP on SNP-chip for genome-wide analysis of human histone H4 hyperacetylation. *BMC Genomics.* 2007 Sep 14;8:322.
4. Rip J, Nierman MC, Sierts JA, Petersen W, Van den Oever K, Van Raalte D, Ross CJ, Hayden MR, Bakker AC, Dijkhuizen P, Hermens WT, Twisk J, Stroes E, Kastelein JJ, Kuivenhoven JA, Meulenberg JM. Human Gene Ther. 2005 Nov; 16(11): 1276-86.
5. Saitoh A, Haas RH, Naviaux RK, Salva NG, Wong JK, Spector SA. Impact of nucleoside reverse transcriptase inhibitors on mitochondrial DNA and RNA in human skeletal muscle cells. *Antimicrob Agents Chemother.* 2008 Aug;52(8):2825-30.
6. Stern MM, Myers RL, Hammam N, Stern KA, Eberli D, Kritchevsky SB, Soker S, Van Dyke M. The influence of extracellular matrix derived from skeletal muscle tissue on the proliferation and differentiation of myogenic progenitor cells ex vivo. *Biomaterials.* 2009 Apr;30(12):2393-9.

7. Thorrez L, Shansky J, Wang L, Fast L, VandenDriessche T, Chuah M, Mooney D, Vandenburgh H. Growth, differentiation, transplantation and survival of human skeletal myofibers on biodegradable scaffolds. *Biomaterials*. 2008 Jan;29(1):75-84.

\* References not specifically citing the use of Lonza cells, media, or reagents in their research.

+ Denotes sections containing only the articles published within the last ten years.